

## SEARCH REQUEST FORM

## Scientific and Technical Information Center

Requester's Full Name: Rob Rabago Examiner #: 75055 Date: 9/10/02  
 Art Unit: 1713 Phone Number 308-4347 Serial Number: 09/776,984  
 Mail Box and Bldg/Room Location: CP3 8B32 Results Format Preferred (circle): PAPER DISK E-MAIL  
CP3 9A09

If more than one search is submitted, please prioritize searches in order of need.

\*\*\*\*\*

Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched.

Include the elected species or structures; keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc, if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.

Title of Invention: Catalyst compositions for the polymerization of olefins

Inventors (please provide full names): James Allen Ponasik, Jason Patrick McDevitt,  
Christopher Moore Killian, Peter Border Mackenzie, Leslie Shane Moody

Earliest Priority Filing Date: 3/13/97

\*For Sequence Searches Only\* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.

The process of claim 1


BEST AVAILABLE COPY

STAFF USE ONLY		Type of Search	Vendors and cost where applicable
Searcher: <u>K. Fuller</u>		NA Sequence (#)	STN <u>✓</u>
Searcher Phone #:		AA Sequence (#)	Dialog
Searcher Location:		Structure (#)	<u>6</u> Questel/Orbit
Date Searcher Picked Up:		Bibliographic	Dr. Link
Date Completed: <u>9/13/02</u>		Litigation	Lexis/Nexis
Searcher Prep & Review Time: <u>20</u>		Fulltext	Sequence Systems
Clerical Prep Time:		Patent Family	WWW/Internet
Online Time: <u>55</u>		Other	Other (specify)

# EIC1700

## Search Results

### Feedback Form (Optional)



Scientific & Technical Information Center

The search results generated for your recent request are attached. If you have any questions or comments (compliments or complaints) about the scope or the results of the search, please contact *the EIC searcher* who conducted the search or contact:

Kathleen Fuller, Team Leader, 308-4290, CP3/4 3D62

#### *Voluntary Results Feedback Form*

➤ *I am an examiner in Workgroup:*  Example: **1713**

➤ *Relevant prior art found, search results used as follows:*

- 102 rejection
- 103 rejection
- Cited as being of interest.
- Helped examiner better understand the invention.
- Helped examiner better understand the state of the art in their technology.

*Types of relevant prior art found:*

- Foreign Patent(s)
- Non-Patent Literature  
(journal articles, conference proceedings, new product announcements etc.)

➤ *Relevant prior art not found:*

- Results verified the lack of relevant prior art (helped determine patentability).
- Search results were not useful in determining patentability or understanding the invention.

**Other Comments:**

---

Drop off completed forms in CP3/4 - 3D62 .

=> FILE REG  
FILE 'REGISTRY' ENTERED AT 12:36:39 ON 13 SEP 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 12 SEP 2002 HIGHEST RN 450335-18-9  
DICTIONARY FILE UPDATES: 12 SEP 2002 HIGHEST RN 450335-18-9

TSCA INFORMATION NOW CURRENT THROUGH MAY 20, 2002

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Calculated physical property data is now available. See HELP PROPERTIES  
for more information. See STNote 27, Searching Properties in the CAS  
Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=> FILE HCAPLUS  
FILE 'HCAPLUS' ENTERED AT 12:36:44 ON 13 SEP 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is  
held by the publishers listed in the PUBLISHER (PB) field (available  
for records published or updated in Chemical Abstracts after December  
26, 1996), unless otherwise indicated in the original publications.  
The CA Lexicon is the copyrighted intellectual property of the  
the American Chemical Society and is provided to assist you in searching  
databases on STN. Any dissemination, distribution, copying, or storing  
of this information, without the prior written consent of CAS, is  
strictly prohibited.

FILE COVERS 1907 - 13 Sep 2002 VOL 137 ISS 12  
FILE LAST UPDATED: 12 Sep 2002 (20020912/ED)

This file contains CAS Registry Numbers for easy and accurate  
substance identification.

CAS roles have been modified effective December 16, 2001. Please  
check your SDI profiles to see if they need to be revised. For  
information on CAS roles, enter HELP ROLES at an arrow prompt or use  
the CAS Roles thesaurus (/RL field) in this file.

=> D QUE L31  
L5 STR  
M 5 N->C->G1->N  
1 2 3 4

VAR G1=C/N  
NODE ATTRIBUTES:  
NSPEC IS RC AT 1  
NSPEC IS RC AT 2  
NSPEC IS RC AT 4

225,604 structures  
from this  
query


NSPEC IS RC AT 5  
DEFAULT MLEVEL IS ATOM  
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:  
RING(S) ARE ISOLATED OR EMBEDDED  
NUMBER OF NODES IS 5

STEREO ATTRIBUTES: NONE

L7 SCR 1993  
L8 SCR 1965

L10 225604 SEA FILE=REGISTRY SSS FUL L5 AND L7 AND L8  
L21 STR



*5 subset searches with  
3 more exact  
structure query*


NODE ATTRIBUTES:

CONNECT IS M3 R AT 2  
CONNECT IS M3 R AT 3  
DEFAULT MLEVEL IS ATOM  
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
NUMBER OF NODES IS 7

STEREO ATTRIBUTES: NONE  
L22 STR



NODE ATTRIBUTES:


CONNECT IS M3 R AT 2  
CONNECT IS M3 R AT 3  
DEFAULT MLEVEL IS ATOM  
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
NUMBER OF NODES IS 7

STEREO ATTRIBUTES: NONE  
L23 STR

*next page*



VAR G1=CB/AK

NODE ATTRIBUTES:

NSPEC IS RC AT 7  
 DEFAULT MLEVEL IS ATOM  
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
 NUMBER OF NODES IS 9

STEREO ATTRIBUTES: NONE

L25 22626 SEA FILE=REGISTRY SUB=L10 SSS FUL (L21 OR L22 OR L23)  
 L27 8931 SEA FILE=HCAPLUS ABB=ON L25  
 L29 392 SEA FILE=HCAPLUS ABB=ON L27(L)LIGAND#  
 L30 44405 SEA FILE=HCAPLUS ABB=ON POLYMERI?(4A) (ETHYLENE OR PROPYLENE  
     OR BUTYLENE OR OLEFIN#)  
 L31 17 SEA FILE=HCAPLUS ABB=ON L29 AND L30

=&gt; D L31 ALL 1-17 HITSTR

L31 ANSWER 1 OF 17 HCAPLUS COPYRIGHT 2002 ACS

AN 2001:609863 HCAPLUS

DN 135:181095

TI Manufacture of supported olefin polymerization catalysts containing transition metal-nitrogen tridentate ligand complex

IN Payne, Mark John

PA BP Chemicals Ltd., UK

SO Jpn. Kokai Tokkyo Koho, 11 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM C08F004-70

ICS C08F010-00

CC 35-3 (Chemistry of Synthetic High Polymers)

Section cross-reference(s): 67

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2001226424	A2	20010821	JP 2001-34799	20010213
	EP 1125952	A1	20010822	EP 2001-300584	20010123
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	US 2001044374	A1	20011122	US 2001-772880	20010131
PRAI	GB 2000-3356	A	20000214		
OS	MARPAT 135:181095				
AB	The catalysts, which are cheap and have uniform activity, are prep'd. by steps of: (1) contacting a support contg. 1-10% water, e.g., silica, alumina, aluminosilicate or crosslinked polystyrene or poly(vinyl alc.), with trialkylaluminum compd., and (2) contacting the product from step 1				

22,626 structures  
 from subset  
 search

with a transition metal-nitrogen tridentate ligand complex. Thus, reacting 2.0 g 2,6-diacetylpyridinebis(2,4,6-trimethylanil) with 0.638 g iron dichloride in n-butanol at 80.degree. for 60 min and at room temp. for 16 h gave 2.56 g 2,6-diacetylpyridinebis(2,4,6-trimethylanil)iron dichloride, 0.0262 g of which (in dried MePh) was mixed with silica/MAO slurry prep'd. in situ from 2 g ES 70X silica and 0.00494 mol trimethylaluminum to give a title catalyst with 0.12% Fe and 12.5% MAO, which was used for **ethylene** slurry **polymn.** with addnl. 3 mL 1 M triisobutylaluminum to give polyethylene with catalyst activity 5778 g/mmol-Fe/h.cntdot.Bar.

ST supported transition metal nitrogen tridentate ligand complex manuf; iron complex olefin ethylene polymn catalyst manuf

IT Polymerization catalysts  
(coordination; manuf. of supported **olefin polymn.**  
catalysts contg. transition metal-nitrogen tridentate ligand complex)

IT Transition metal complexes  
RL: CAT (Catalyst use); IMF (Industrial manufacture); PREP (Preparation);  
USES (Uses)  
(nitrogen heterocyclic; manuf. of supported **olefin polymn.**  
catalysts contg. transition metal-nitrogen tridentate ligand complex)

IT Heterocyclic compounds  
RL: CAT (Catalyst use); IMF (Industrial manufacture); PREP (Preparation);  
USES (Uses)  
(nitrogen, transition metal complexes; manuf. of supported  
**olefin polymn.** catalysts contg. transition metal-nitrogen tridentate ligand complex)

IT Polymerization catalysts  
(supported; manuf. of supported **olefin polymn.**  
catalysts contg. transition metal-nitrogen tridentate ligand complex)

IT 7631-86-9, ES 70X, reactions  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(ES 70X, catalyst support; in manuf. of supported **olefin polymn.** catalysts contg. iron-nitrogen tridentate ligand complex)

IT 210155-39-8P  
RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)  
(catalyst intermediate; in manuf. of supported **olefin polymn.** catalysts contg. iron-nitrogen tridentate ligand complex)

IT 100-99-2, Triisobutylaluminum, uses  
RL: CAT (Catalyst use); USES (Uses)  
(cocatalyst; manuf. of supported **olefin polymn.**  
catalysts contg. iron-nitrogen tridentate ligand complex)

IT 308359-84-4P  
RL: CAT (Catalyst use); IMF (Industrial manufacture); PREP (Preparation);  
USES (Uses)  
(manuf. of supported **olefin polymn.** catalysts  
contg. iron-nitrogen tridentate ligand complex)

IT 9002-88-4P, Polyethylene  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(manuf. of supported **olefin polymn.** catalysts  
contg. iron-nitrogen tridentate ligand complex)

IT 207129-93-9 207129-95-1 207129-96-2  
210537-35-2 210768-87-9 221391-06-6 221391-08-8\*\*  
\* \*\*\*221391-12-4 221391-13-5 221391-15-7  
261787-81-9 308359-85-5 308359-86-6  
355118-93-3


IT 88-05-1, 2,4,6-Trimethylaniline 1129-30-2, 2,6-Diacetylpyridine  
 7758-94-3, Iron dichloride  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reactant for catalyst; in manuf. of supported **olefin**  
**polymn.** catalysts contg. iron-nitrogen tridentate ligand  
 complex)

IT 75-24-1, Trimethylaluminum 97-93-8, Triethylaluminum, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reactant for cocatalyst; in manuf. of supported **olefin**  
**polymn.** catalysts contg. iron-nitrogen tridentate ligand  
 complex)

IT 308359-84-4P  
 RL: CAT (Catalyst use); IMF (Industrial manufacture); PREP (Preparation);  
 USES (Uses)  
 (manuf. of supported **olefin** **polymn.** catalysts  
 contg. iron-nitrogen tridentate ligand complex)

RN 308359-84-4 HCPLUS


CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,4,6-trimethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



IT 207129-93-9 207129-95-1 207129-96-2  
 210537-35-2 210768-87-9 221391-08-8  
 221391-12-4 221391-13-5 221391-15-7  
 261787-81-9 308359-85-5 308359-86-6  
 355118-93-3  
 RL: CAT (Catalyst use); USES (Uses)  
 (manuf. of supported **olefin** **polymn.** catalysts  
 contg. transition metal-nitrogen tridentate **ligand** complex)


RN 207129-93-9 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-dimethylbenzenamine-.kappa.N]-, (SP-5-13)- (9CI) (CA INDEX NAME)




RN 207129-95-1 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,4-dimethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



RN 207129-96-2 HCAPLUS


CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethylidyne]bis[2,6-dimethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



RN 210537-35-2 HCAPLUS


KATHLEEN FULLER EIC 1700/LAW LIBRARY 308-4290

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)bis(ethylidynenitrilo-.kappa.N)]bis[2-methylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)




RN 210768-87-9 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliyne]bis[2-(1,1-dimethylethyl)benzenamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)




RN 221391-08-8 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliyne]bis[2,3-dimethylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)




RN 221391-12-4 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethylidyne]bis[2,6-diisopropylbenzylamine-.kappa.N]- (9CI) (CA INDEX NAME)



RN 221391-13-5 HCPLUS


CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethylidyne]bis[2,6-bis(1-methylethyl)benzylamine-.kappa.N]- (9CI) (CA INDEX NAME)



RN 221391-15-7 HCPLUS


CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethylidyne]bis[1-

naphthalenamine-.kappa.N]]- (9CI) (CA INDEX NAME)




RN 261787-81-9 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[4-(1,1-dimethylethyl)-2,6-dimethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)




RN 308359-85-5 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]- (9CI) (CA INDEX NAME)




RN 308359-86-6 HCAPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)



RN 355118-93-3 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)bis(phenylmethylidyne)]bis[benzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)



AN 2001:584639 HCAPLUS  
 DN 135:304175  
 TI **Polymerization of ethylene** catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand: 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe3, MAO, AlMe3/B(C6F5)3 and AlMe3/CPh3B(C6F5)4  
 AU Talsi, Evgenii P.; Babushkin, Dmitrii E.; Semikolenova, Nina V.; Zudin, Vladimir N.; Panchenko, Valentina N.; Zakharov, Vladimir A.  
 CS Boreskov Institute of Catalysis, Novosibirsk, 630090, Russia  
 SO Macromolecular Chemistry and Physics (2001), 202(10), 2046-2051  
 CODEN: MCHPES; ISSN: 1022-1352

---

PB Wiley-VCH Verlag GmbH  
 DT Journal  
 LA English  
 CC 35-3 (Chemistry of Synthetic High Polymers)  
 Section cross-reference(s): 29, 67  
 AB 1H and 2H NMR spectroscopic monitoring of ferrous species formed via interaction of 2,6-bis[1-(2,6-dimethylphenylimino)ethyl]pyridineiron(II) chloride (1) with AlMe3, MAO, AlMe3/B(C6F5)3 and AlMe3/CPh3 (C6F5)4 is reported. At interaction of 1 with MAO in toluene soln., the new stable heterodinuclear neutral complexes with proposed structures LFe(II)(Cl)(.mu.-Me)2AlMe2 and LFe(II)(Me)(.mu.-Me)2AlMe2 are formed (L is initial tridentate ligand). Complex LFe(II)(Cl)(.mu.-Me)2AlMe2 predominates at low Al/Fe ratios (less than 50), while LFe(II)(Me)(.mu.-Me)2AlMe2 at high Al/Fe ratios (more than 500). Complex assigned to LFe(II)(Me)(.mu.-Me)2AlMe2 can be prep'd. via interaction of 1 with AlMe3. Activation of LFe(II)(Me)(.mu.-Me)2AlMe2 by B(C6F5)3 and CPh3B(C6F5)4 gives rise to formation of new complexes with proposed structures [LFe(.mu.-Me)2AlMe2]+[MeB(C6F5)3]- and [LFe(.mu.-Me)2AlMe2]+[B(C6F5)4]-. Unexpectedly, the activity at **ethylene polymn.** was even higher for 1/AlMe3 than for 1/MAO catalytic system. The co-catalytic activity of MAO towards 1 dramatically decreased with the diminishing of AlMe3 content in the compon. of MAO. Activity of the catalyst 1/AlMe3 and the mol. structure of polyethylene produced do not change noticeably at the addn. of B(C6F5)3 to 1/AlMe3. These data allow to suggest, that active species of 1/AlMe3 and 1/MAO systems are neutral methylated ferrous complexes but not cationic intermediates. Probably, complex LFe(II)(Me)(.mu.-Me)2 AlMe2 is the closest precursor of these active species.

ST iron complex bisiminepyridyl ligand catalyst **ethylene polymn.**; aluminum cocatalyst iron complex polymn catalyst interaction; boron compd cocatalyst iron complex polymn catalyst interaction

IT Aluminoxanes  
 RL: CAT (Catalyst use); RCT (Reactant); RACT (Reactant or reagent); USES (Uses)  
 (Me, cocatalyst; polymn. of **ethylene** catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe3, MAO, AlMe3/B(C6F5)3 and AlMe3/CPh3B(C6F5)4)

IT **Polymerization catalysts**  
 (polymn. of **ethylene** catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe3, MAO, AlMe3/B(C6F5)3 and AlMe3/CPh3B(C6F5)4)

IT 75-24-1, Trimethylaluminum 1109-15-5, Perfluorotriphenylboron 136040-19-2, Triphenylmethyl tetrakis(pentafluorophenyl)borate  
 RL: CAT (Catalyst use); RCT (Reactant); RACT (Reactant or reagent); USES (Uses)  
 (cocatalyst; polymn. of **ethylene** catalyzed by iron

complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe<sub>3</sub>, MAO, AlMe<sub>3</sub>/B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and AlMe<sub>3</sub>/CPh3B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>)

IT 207129-93-9P  
 RL: CAT (Catalyst use); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)  
 (polymn. of ethylene catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe<sub>3</sub>, MAO, AlMe<sub>3</sub>/B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and AlMe<sub>3</sub>/CPh3B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>)

IT 350679-35-5P 350679-36-6P 350679-39-9P 350679-40-2P  
 RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation); USES (Uses)  
 (polymn. of ethylene catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe<sub>3</sub>, MAO, AlMe<sub>3</sub>/B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and AlMe<sub>3</sub>/CPh3B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>)

IT 87-62-7, 2,6-Dimethylaniline 1129-30-2, 2,6-Diacetylpyridine  
 7758-94-3, Ferrous chloride  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (polymn. of ethylene catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe<sub>3</sub>, MAO, AlMe<sub>3</sub>/B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and AlMe<sub>3</sub>/CPh3B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>)

IT 204203-16-7P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (polymn. of ethylene catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe<sub>3</sub>, MAO, AlMe<sub>3</sub>/B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and AlMe<sub>3</sub>/CPh3B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>)


IT 9002-88-4P, Polyethylene  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (polymn. of ethylene catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe<sub>3</sub>, MAO, AlMe<sub>3</sub>/B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and AlMe<sub>3</sub>/CPh3B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>)

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE

(1) Bochmann, M; Angew Chem, Int Ed Engl 1994, V33, P1634  
 (2) Brintzinger, H; Angew Chem, Int Ed Engl 1995, V34, P1143 HCPLUS  
 (3) Britovsek, G; Angew Chem, Int Ed Engl 1999, V38, P428 HCPLUS  
 (4) Britovsek, G; Chem Commun 1998, P849  
 (5) Britovsek, G; J Am Chem Soc 1999, V121, P8728 HCPLUS  
 (6) Hill, D; J Am Chem Soc 1988, V110, P1651  
 (7) Small, B; J Am Chem Soc 1998, V120, P4049 HCPLUS  
 (8) Yang, X; J Am Chem Soc 1991, V113, P3623 HCPLUS

IT 207129-93-9P  
 RL: CAT (Catalyst use); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)  
 (polymn. of ethylene catalyzed by iron complex bearing 2,6-bis(imine)pyridyl ligand and 1H and 2H NMR monitoring of ferrous species formed via catalyst activation with AlMe<sub>3</sub>, MAO, AlMe<sub>3</sub>/B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and AlMe<sub>3</sub>/CPh3B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>)

RN 207129-93-9 HCPLUS  
 CN Iron, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-dimethylbenzenamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)



L31 ANSWER 3 OF 17 HCAPLUS COPYRIGHT 2002 ACS  
 AN 2001:302382 HCAPLUS  
 DN 134:326893  
 TI Tridentate ligands based on bis(iminomethyl)pyridine derivatives  
 IN Bennett, Alison  
 PA USA  
 SO U.S. Pat. Appl. Publ., 17 pp., Cont. -in-part of U. S. Ser. No. 273,409.  
 CODEN: USXXCO


DT Patent  
 LA English  
 IC ICM C07D211-70  
 ICS C08F220-10

NCL 526329000

CC 35-3 (Chemistry of Synthetic High Polymers)  
 Section cross-reference(s): 67

FAN.CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2001000519	A1	20010426	US 2000-729305	20001204
	US 6423848	B2	20020723		
	US 5955555	A	19990921	US 1997-991372	19971216
	EP 1127897	A2	20010829	EP 2001-200886	19971216
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI				
	US 6214761	B1	20010410	US 1999-273409	19990322
PRAI	US 1996-33656P	P	19961217		
	US 1997-991372	A2	19971216		
	US 1999-273409	A2	19990322		
	EP 1997-953340	A3	19971216		
OS	MARPAT 134:326893				
GI					



AB Title derivs. I [R1-3, R9-16 = H, (substituted) hydrocarbyl, or inert functional group; R4, R5 = H or (substituted) hydrocarbyl, R8, R12, R13, R17 = (substituted) hydrocarbyl or inert functional group, any 2 of R8-17 that are vicinal to one another taken together may form a ring] are useful for as Co or Fe complexes for catalysts in **polymn.** of **olefins**.

ST bisiminomethylpyridine deriv cobalt complex catalyst **polymn** **olefin**; iron bisiminomethylpyridine deriv complex catalyst **polymn** **olefin**

IT Aluminoxanes  
RL: CAT (Catalyst use); USES (Uses)  
(Me, cocatalyst; tridentate ligands based on bis(iminomethyl)pyridine derivs. for cobalt or iron complexes for catalysts for **polymn** . of **olefins**)

IT Polymerization catalysts  
(tridentate ligands based on bis(iminomethyl)pyridine derivs. for cobalt or iron complexes for catalysts for **polymn.** of **olefins**)

IT 1271-19-8, Titanocene dichloride 54039-38-2 75171-00-5 75180-85-7  
85959-83-7 112243-79-5 119821-97-5 130638-44-7 132510-07-7  
135072-61-6 135539-49-0 148799-37-5 148799-45-5 149342-08-5  
156367-60-1 191155-99-4 245360-94-5 245361-10-8 245361-12-0  
245361-15-3 245361-17-5  
RL: CAT (Catalyst use); USES (Uses)  
(catalyst; tridentate ligands based on bis(iminomethyl)pyridine derivs. for cobalt or iron complexes for catalysts for **polymn.** of **olefins**)

IT 210537-35-2 308359-84-4 308359-85-5  
308359-86-6  
RL: CAT (Catalyst use); USES (Uses)  
(tridentate **ligands** based on bis(iminomethyl)pyridine derivs. for cobalt or iron complexes for catalysts for **polymn.** of **olefins**)

IT 9002-88-4P, Polyethylene 9003-07-0P, Polypropylene  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(tridentate ligands based on bis(iminomethyl)pyridine derivs. for cobalt or iron complexes for catalysts for **polymn.** of


olefins)

IT 204203-14-5 210155-39-8 210155-40-1 210155-55-8 335657-86-8  
 RL: MSC (Miscellaneous)  
 (tridentate ligands based on bis(iminomethyl)pyridine derivs. for cobalt or iron complexes for catalysts for polymn. of olefins)


IT 210537-35-2 308359-84-4 308359-85-5  
 308359-86-6  
 RL: CAT (Catalyst use); USES (Uses)  
 (tridentate ligands based on bis(iminomethyl)pyridine derivs. for cobalt or iron complexes for catalysts for polymn. of olefins)

RN 210537-35-2 HCAPLUS


CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)bis(ethylidynenitrilo-.kappa.N)]bis[2-methylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



RN 308359-84-4 HCAPLUS  
 CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,4,6-trimethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)




RN 308359-85-5 HCAPLUS  
 CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



RN 308359-86-6 HCAPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



L31 ANSWER 4 OF 17 HCAPLUS COPYRIGHT 2002 ACS

AN 2001:246353 HCAPLUS

DN 135:77129

TI FI Catalysts: super active new **ethylene** polymerization catalysts

AU Matsui, S.; Fujita, T.

CS Materials Science Laboratory, 530-32 Nagaura, Mitsui Chemicals, Inc., Chiba, Sodegaura-City, Japan

SO Catalysis Today (2001), 66(1), 63-73  
CODEN: CATTEA; ISSN: 0920-5861

PB Elsevier Science B.V.

DT Journal

LA English

CC 35-3 (Chemistry of Synthetic High Polymers)  
Section cross-reference(s): 78

AB Based on "ligand-oriented catalyst design", group 4 transition metal complexes having two phenoxy-imine ligands, FI Catalysts, were synthesized and investigated as **olefin** polymer catalysts using **ethylene** at atm. pressure. As a result, these complexes were

found to exhibit very high activity using MAO as a cocatalyst. Among them, zirconium complexes displayed the highest activity with moderate (M<sub>v</sub>: 0.8.times.10<sup>4</sup>) to very high (M<sub>v</sub>: 71.6.times.10<sup>4</sup>) mol. wt. ranges. The max. activity exceeded 4 t-PE/(mmol-cat<sup>h</sup>) at 25.degree.C even at atm. pressure, the activity being two orders of magnitude larger than that exhibited by Cp<sub>2</sub>ZrCl<sub>2</sub>. Alternatively, by using borate/iBu<sub>3</sub>Al as a cocatalyst, a zirconium FI Catalyst produced exceptionally high mol. wt. polyethylene (M<sub>v</sub>: 505.times.10<sup>4</sup>) displaying considerable activity. As far as we know, this is one of the highest mol. wt. values obtained from homogeneous polymn. catalysts. These results indicate that FI Catalysts possess very high potential as olefin polymn. catalysts.

ST group IV transition metal complex phenoxyimine ligand polymn catalyst; **ethylene polymn** transition metal complex phenoxyimine ligand catalyst

IT Aluminoxanes

RL: CAT (Catalyst use); USES (Uses)  
(Me; group IV transition metal complexes having two phenoxy-imine ligands as super active **ethylene polymn.** catalysts)

IT Polymerization catalysts

(group IV transition metal complexes having two phenoxy-imine ligands as super active **ethylene polymn.** catalysts)

IT 100-99-2, Tris(isobutyl) aluminum, uses 136040-19-2, Triphenylcarbonium tetrakis(pentafluorophenyl)borate 215050-65-0 215050-67-2  
215051-00-6 215051-04-0 215051-12-0 215051-20-0 **215594-94-8**  
261375-85-3 264626-79-1 279218-85-8 279218-86-9 **347850-82-2**  
347850-83-3

RL: CAT (Catalyst use); USES (Uses)  
(group IV transition metal complexes having two phenoxy-imine ligands as super active **ethylene polymn.** catalysts)

IT 9002-88-4P, Polyethylene

RL: SPN (Synthetic preparation); PREP (Preparation)  
(group IV transition metal complexes having two phenoxy-imine ligands as super active **ethylene polymn.** catalysts)

RE.CNT 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Alt, H; J Organomet Chem 1994, V472, P113 HCPLUS
- (2) Anon; Korehasugoi! Kagaku no Sekai Kiroku-shu 1999, P111
- (3) Bei, X; Organometallics 1997, V16, P3282 HCPLUS
- (4) Biritovsek, G; Angew Chem Int Ed Engl 1999, V38, P428
- (5) Bochmann, M; J Chem Soc, Dalton Trans 1996, P255 HCPLUS
- (6) Brintzinger, H; Angew Chem Int Ed Engl 1995, V34, P1143 HCPLUS
- (7) Britovsek, G; Chem Commun 1998, P849
- (8) Chiang, R; J Polym Sci 1959, V36, P91 HCPLUS
- (9) Deck, P; J Am Chem Soc 1998, V120, P1772 HCPLUS
- (10) Deng, L; Organometallics 1998, V17, P3240 HCPLUS
- (11) Flores, J; Organometallics 1995, V14, P1827 HCPLUS
- (12) Fonseca Gueerra, C; Theor Chem Acc 1998, V99, P391
- (13) Fujita, T; EP 0874005 1998 HCPLUS
- (14) Fujita, T; Proceedings of the MetCon'99 on Polymer in Transition
- (15) Hakala, K; Macromol Chem Rapid Commun 1997, V18, P635 HCPLUS
- (16) Johnson, L; J Am Chem Soc 1995, V117, P6414 HCPLUS
- (17) Johnson, L; J Am Chem Soc 1996, V118, P267 HCPLUS
- (18) Jones, D; J Chem Soc, Dalton Trans 1998, P255 HCPLUS
- (19) Kaim, W; Angew Chem Int Ed Engl 1978, V17, P466
- (20) Kaim, W; Angew Chem Int Ed Engl 1981, V20, P116
- (21) Kashiwa, N; US 3642746 1968 HCPLUS
- (22) Killian, C; J Am Chem Soc 1996, V118, P11664 HCPLUS
- (23) Kim, I; Organometallics 1997, V16, P3314 HCPLUS

(24) Klabunde, U; J Mol Catal 1987, V41, P123 HCPLUS  
 (25) Klabunde, U; J Mol Catal 1987, V41, P123 HCPLUS  
 (26) Linden, A; J Am Chem Soc 1995, V117, P3008  
 (27) Mackenzie, P; Proceedings of the MetCon'99 on Polymer in Transition  
 (28) Matilainen, L; J Chem Soc, Dalton Trans 1996, P219 HCPLUS  
 (29) Matsui, S; Chem Lett 1999, P1065 HCPLUS  
 (30) Matsui, S; Chem Lett 1999, P1163  
 (31) Matsui, S; Chem Lett 2000, P554 HCPLUS  
 (32) Mecking, S; J Am Chem Soc 1998, V120, P888 HCPLUS  
 (33) Miyatake, T; Makromol Chem Rapid Commun 1989, V10, P349 HCPLUS  
 (34) Mohring, V; Angew Chem Int Ed Engl 1985, V24, P1001


---

(35) Saeki, K; JP 7319601 1973 HCPLUS  
 (36) Saeki, K; JP 7322682 1973 HCPLUS  
 (37) Saito, J; JP 324709 A 1998  
 (38) Schattenmann, F; Organometallics 1998, V17, P989 HCPLUS  
 (39) Scollard, J; Makromolekules 1996, V29, P5241 HCPLUS  
 (40) Scollard, J; Organometallics 1997, V16, P1810 HCPLUS  
 (41) Scollard, J; Organometallics 1997, V16, P4415 HCPLUS  
 (42) Sinn, H; Adv Organomet Chem 1980, V18, P99 HCPLUS  
 (43) Sinn, H; Angew Chem Int Ed Engl 1980, V19, P390  
 (44) Small, B; J Am Chem Soc 1998, V120, P4049 HCPLUS  
 (45) Spaleck, W; Organometallics 1994, V13, P954 HCPLUS  
 (46) Stephan, D; Organometallics 1999, V18, P2046 HCPLUS  
 (47) Suzuki, N; Macromolecules 2000, V33, P754 HCPLUS  
 (48) Tjaden, E; Organometallics 1995, V14, P371 HCPLUS  
 (49) Tsukahara, T; Organometallics 1997, V16, P3303 HCPLUS  
 (50) Vollmerhaus, R; Organometallics 1999, V18, P2731 HCPLUS  
 (51) Wang, C; Organometallics 1998, V17, P3149 HCPLUS  
 (52) Younkin, T; Science 2000, V287, P460 HCPLUS  
 (53) Ziegler, K; Angew Chem 1955, V67, P424  
 (54) Ziegler, K; Angew Chem 1955, V67, P541 HCPLUS

IT 215594-94-8 347850-82-2


RL: CAT (Catalyst use); USES (Uses)  
 (group IV transition metal complexes having two phenoxy-imine  
 ligands as super active ethylene polymn.  
 catalysts)

RN 215594-94-8 HCPLUS

CN Nickel, [2,6-bis(1-methylethyl)-N-[(2-pyridinyl-  
 .kappa.N)methylene]benzenamine-.kappa.N]dibromo- (9CI) (CA INDEX NAME)

RN 347850-82-2 HCPLUS

CN Cobalt, dibromo[2-(phenoxy-.kappa.O)-N-[(2-pyridinyl-  
 .kappa.N)methylene]benzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



L31 ANSWER 5 OF 17 HCAPLUS COPYRIGHT 2002 ACS  
 AN 2001:178430 HCAPLUS  
 DN 134:208327  
 TI Transition metal complexes of imino nitrogen-containing heterocycle  
 ligands as catalysts for the polymerization of olefins  
 IN Ponasik, James Allen, Jr.; McDevitt, Jason Patrick; Killian, Christopher  
 Moore; MacKenzie, Peter Borden; Moody, Leslie Shane; Lavoie, Gino Georges  
 PA Eastman Chemical Company, USA  
 SO U.S., 27 pp., Cont.-in-part of U.S. Ser. No. 28,315 *applicante*  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 IC ICM B01J023-38  
 ICS B01J023-40; B01J023-74; B01J023-75; B01J023-755  
 NCL 502162000  
 CC 35-3 (Chemistry of Synthetic High Polymers)  
 Section cross-reference(s): 29  
 FAN.CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6200925	B1	20010313	US 1998-222614	19981229
	US 2002035030	A1	20020321	US 2001-776984	20010205
	US 2001025007	A1	20010927	US 2001-780093	20010209
	US 6372682	B2	20020416		
PRAI	US 1997-40754P	P	19970313		
	US 1997-44691P	P	19970418		
	US 1997-45337P	P	19970501		
	US 1997-45357P	P	19970502		
	US 1997-45358P	P	19970502		
	US 1997-45697P	P	19970506		
	US 1998-28315	A2	19980224		
	US 1998-222614	A3	19981229		

OS MARPAT 134:208327  
 AB The title catalyst systems, useful in the polymerization of  
 olefins, comprise a Group 8-10 transition metal component and a  
 ligand component comprising a N and/or functional groups comprising a N,  
 generally in the form of an imine functional group. In certain  
 embodiments, the ligand component may further comprise a phosphorus atom.  
 Preferred ligand components are bidentate (bind to the transition metal at  
 two or more sites) and include a nitrogen-transition metal bond. The  
 transition metal-ligand complex is generally cationic and associated with a  
 weakly coordinating anion. The catalyst system may be supported or  
 comprises a Lewis or Bronsted acid complexed with the ligand component of  
 the transition metal-ligand complex, e.g. Me aluminoxane or B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>.  
 ST nickel complex olefin polymer catalysts  
 IT Aluminoxanes  
 RL: CAT (Catalyst use); USES (Uses)  
 (Me; Transition metal complexes of imino or nitrogen-contg. heterocycle

ligands as catalysts for the **polymn. of olefins**)

IT Polyolefins  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn. of olefins**)

IT Ligands  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn. of olefins**)

IT Polymerization catalysts  
 (coordination; Transition metal complexes of imino or nitrogen-contg.  
 heterocycle ligands as catalysts for the **polymn. of olefins**)

IT 1109-15-5, Tris(pentafluorophenyl)borane 121281-53-6 **328565-01-1**  
 RL: CAT (Catalyst use); USES (Uses)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn. of olefins**)

IT 213384-29-3P 213384-31-7P 213384-33-9P  
 213384-35-1P 213384-37-3P 213384-39-5P  
 213384-41-9P 213384-43-1P 213384-45-3P  
 213384-47-5P 213384-49-7P 213384-51-1P  
 213384-53-3P 213384-55-5P 213384-57-7P  
 328565-00-0P 328565-02-2P 328565-03-3P  
 328565-04-4P 328565-05-5P 328565-06-6P  
 328565-08-8P 328565-10-2P  
 RL: CAT (Catalyst use); IMF (Industrial manufacture); PREP (Preparation);  
 USES (Uses)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn. of olefins**)

IT 1295-35-8  
 RL: CAT (Catalyst use); RCT (Reactant); RACT (Reactant or reagent); USES  
 (Uses)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn. of olefins**)

IT 9002-88-4P, Polyethylene 25038-76-0P, Norbornene, homopolymer  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn. of olefins**)

IT 121-44-8, Triethylamine, reactions 271-44-3, Indazole 273-34-7,  
 1H-1,2,3-Triazolo[4,5-b]pyridine 288-13-1, Pyrazole 288-36-8,  
 1,2,3-Triazole 288-88-0, 1H-1,2,4-Triazole 463-71-8, Thiophosgene  
 1453-58-3 1989-53-3, 2,6-Dimethoxybenzoyl chloride 5401-94-5,  
 5-Nitroindazole 7550-45-0, Titanium tetrachloride, reactions  
 14190-59-1, 2-Thiazolecarboxylic acid 18039-42-4, 5-Phenyl-1H-tetrazole  
 23814-12-2, 1H-Benzotriazole-5-carboxylic acid 24295-03-2,  
 2-Acetylthiazole 24544-04-5 28923-39-9 59387-01-8 64594-47-4  
 157894-08-1 213182-96-8 213182-97-9 213182-98-0 213183-09-6  
 303117-48-8 328565-11-3 328565-12-4 328565-13-5 328565-14-6  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn. of olefins**)

IT 213182-93-5P 328564-93-8P 328565-09-9P  
 RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (ligand; Transition metal complexes of imino or nitrogen-contg.  
 heterocycle ligands as catalysts for the **polymn. of olefins**)

IT 213182-90-2 213182-92-4 213182-94-6 213182-95-7 213182-99-1

213183-00-7 213183-01-8 213183-02-9 213183-03-0 213183-04-1  
 213183-05-2 213183-06-3 213183-07-4 213183-13-2 213183-16-5  
 328564-94-9 328564-95-0 328564-96-1 328564-97-2 328564-98-3  
 328564-99-4 328565-07-7

RL: RCT (Reactant); RACT (Reactant or reagent)  
 (ligand; Transition metal complexes of imino or nitrogen-contg.  
 heterocycle ligands as catalysts for the **polymn.** of  
**olefins**)

IT 7631-86-9, Silica, uses

RL: CAT (Catalyst use); USES (Uses)  
 (support; Transition metal complexes of imino or nitrogen-contg.  
 heterocycle ligands as catalysts for the **polymn.** of  
**olefins**)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE


- (1) Anon; EP 0748821 1996 HCPLUS
- (2) Anon; WO 9830610 1998 HCPLUS
- (3) Appel, R; Chemische Berichte Jahrg 1973, V106, P3450 HCPLUS
- (4) Brookhart; US 5866663 1999 HCPLUS
- (5) Tamura, K; J Org Chem 1993, V58, P32 HCPLUS

IT 328565-01-1

RL: CAT (Catalyst use); USES (Uses)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn.** of  
**olefins**)

RN 328565-01-1 HCPLUS

CN Nickel, [1-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]phenylmethyl]-1H-pyrazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)




IT 213384-29-3P 213384-31-7P 213384-33-9P  
 213384-35-1P 213384-37-3P 213384-39-5P  
 213384-41-9P 213384-43-1P 213384-45-3P  
 213384-47-5P 213384-49-7P 213384-51-1P  
 213384-53-3P 213384-55-5P 213384-57-7P  
 328565-00-0P 328565-02-2P 328565-03-3P  
 328565-04-4P 328565-05-5P 328565-06-6P  
 328565-08-8P 328565-10-2P

RL: CAT (Catalyst use); IMF (Industrial manufacture); PREP (Preparation);  
 USES (Uses)  
 (Transition metal complexes of imino or nitrogen-contg. heterocycle  
 ligands as catalysts for the **polymn.** of  
**olefins**)


RN 213384-29-3 HCPLUS

CN Nickel, dibromo[1-[[[2,6-dimethylphenyl]imino-.kappa.N]-1-naphthalenylmethyl]-1H-1,2,4-triazole-.kappa.N2]- (9CI) (CA INDEX NAME)




RN 213384-31-7 HCAPLUS

CN Nickel, [1-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]-1-naphthalenylmethyl]-1H-1,2,4-triazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)




RN 213384-33-9 HCAPLUS

CN Nickel, dibromo[1-[[2,6-dimethylphenyl]imino-.kappa.N]phenylmethyl]-1H-1,2,4-triazole-.kappa.N2] - (9CI) (CA INDEX NAME)




RN 213384-35-1 HCAPLUS


CN Nickel, [1-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]phenylmethyl]-1H-1,2,4-triazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)




RN 213384-37-3 HCAPLUS  
CN Nickel, [1-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N](2,6-dimethoxyphenyl)methyl]-1H-1,2,4-triazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)



RN 213384-39-5 HCAPLUS  
CN Nickel, dibromo[1-[(2,6-dimethylphenyl)imino-.kappa.N]-1-naphthalenylmethyl]-3-methyl-1H-pyrazole-.kappa.N2]- (9CI) (CA INDEX NAME)




RN 213384-41-9 HCAPLUS  
CN Nickel, [1-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]-1-naphthalenylmethyl]-3-methyl-1H-pyrazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)




RN 213384-43-1 HCAPLUS

CN Nickel, [1-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]-1-naphthalenylmethyl]-1H-1,2,3-triazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)




RN 213384-45-3 HCAPLUS

CN Nickel, [1-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]phenylmethyl]-5-phenyl-1H-tetrazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)




RN 213384-47-5 HCAPLUS


CN Nickelate(1-), [1-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]phenylmethyl]-1H-benzotriazole-5-carboxylato-.kappa.N2]dibromo-, hydrogen (9CI) (CA INDEX NAME)




RN 213384-49-7 HCAPLUS  
CN Nickel, [3-[[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]phenylmethyl]-3H-1,2,3-triazolo[4,5-b]pyridine-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)



RN 213384-51-1 HCAPLUS  
CN Nickel, [1-[1-[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]-2,2,2-trifluoroethyl]-1H-indazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)




RN 213384-53-3 HCAPLUS  
CN Nickel, [1-[1-[[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]-2,2,2-trifluoroethyl]-5-nitro-1H-indazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)




RN 213384-55-5 HCAPLUS

CN Nickel, [2,6-bis(1-methylethyl)-N-[1-(2-thiazolyl)-.kappa.N3)ethylidene]benzenamine-.kappa.N]dibromo- (9CI) (CA INDEX NAME)




RN 213384-57-7 HCAPLUS

CN Nickel, dibromo[N-[1-(1H-imidazol-2-yl)-.kappa.N3)methylene]-2,6-bis(1-methylethyl)benzenamine-.kappa.N] - (9CI) (CA INDEX NAME)



RN 328565-00-0 HCAPLUS


CN Nickel, [1-[1,1'-biphenyl]-2-yl[(2,6-dimethylphenyl)imino-.kappa.N]methyl]-1H-1,2,4-triazole-.kappa.N2]dibromo- (9CI) (CA INDEX NAME)



RN 328565-02-2 HCAPLUS


CN Nickel, dibromo[1-[(2,6-dimethylphenyl)imino-.kappa.N] [4-

(trifluoromethyl)phenyl]methyl]-1H-1,2,4-triazole-.kappa.N2]- (9CI) (CA INDEX NAME)





RN 328565-03-3 HCAPLUS

CN Nickel, dibromo[1-[(2,6-dimethylphenyl)imino-.kappa.N](4-methoxyphenyl)methyl]-1H-1,2,4-triazole-.kappa.N2]- (9CI) (CA INDEX NAME)




RN 328565-04-4 HCAPLUS

CN Nickel, dibromo[1-[(2,6-dimethylphenyl)imino-.kappa.N](4-nitrophenyl)methyl]-1H-1,2,4-triazole-.kappa.N2]- (9CI) (CA INDEX NAME)




RN 328565-05-5 HCAPLUS

CN Nickel, dibromo[1-[[4-(1,1-dimethylethyl)phenyl][(2,6-dimethylphenyl)imino-.kappa.N]methyl]-1H-1,2,4-triazole-.kappa.N2]- (9CI) (CA INDEX NAME)




RN 328565-06-6 HCAPLUS

CN Nickel, tetrabromo[.mu.-[1,1'-[1,4-phenylenebis[(2,6-dimethylphenyl)imino-.kappa.N]methylene]]bis[1H-pyrazole-.kappa.N2]]di- (9CI) (CA INDEX NAME)




RN 328565-08-8 HCAPLUS

CN Nickel, dibromo[methyl N-[2,6-bis(1-methylethyl)phenyl]-2-thiazolecarboximidothioate-.kappa.N2,.kappa.N3]- (9CI) (CA INDEX NAME)



RN 328565-10-2 HCAPLUS

CN Nickel, dibromo[methyl N-[2,6-bis(1-methylethyl)phenyl]-1H-1,2,4-triazole-1-carboximidothioate-.kappa.N2,.kappa.N3]- (9CI) (CA INDEX NAME)



L31 ANSWER 6 OF 17 HCAPLUS COPYRIGHT 2002 ACS

AN 2001:152696 HCAPLUS

DN 134:208317

TI Bisimidino compounds, their transition metal complexes, and use of the latter as polymerization catalysts


IN Kristen, Marc Olivier; Gonioukh, Andrei; Lilge, Dieter; Lehmann, Stephan; Bildstein, Benno; Amort, Christoph; Malaun, Michael

PA BASF A.-G., Germany

SO PCT Int. Appl., 53 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA German  
 IC ICM C07F015-02  
 ICS C07F015-06; C08F004-70; C08F010-00; C07D213-53; C07D401-14  
 CC 35-3 (Chemistry of Synthetic High Polymers)  
 Section cross-reference(s): 27, 28, 67, 78

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PT	WO 2001014391	A1	20010301	WO 2000-EP7657	20000808
	W: JP, KR, US RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	EP 1204668	A1	20020515	EP 2000-956407	20000808
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
PRAI	DE 1999-19939415	A	19990820		
	WO 2000-EP7657	W	20000808		
OS	MARPAT 134:208317				
GI					



AB The compds. have the general formula I [A = N, O, P, S; R1 = NR5R6; R2 = NR5R6, NR7R8, alkyl, aryl, cycloalkyl; R3, R4 = H, alkyl, aryl, cycloalkyl; NR5R6 forms an (un)substituted 5-, 6- or 7-membered ring, which can be annellated with (un)substituted 5- or 6-membered rings; R7, R8 = alkyl, aryl, cycloalkyl; n = 1, 2]. Thus, MeCOCH2CH2COCHMe2 was condensed with AcHNH2 to give 53% 1-acetamido-2-isopropyl-5-methylpyrrole, which was deacetylated and condensed 2:1 with 2,6-diacetylpyridine to give a diimine, which was complexed with FeCl2. Copolymn. of ethylene with 1-hexene in toluene in the presence of Me aluminoxane and the complex at 30.degree. for 1 h gave a copolymer with catalyst efficiency 980 g/mmol catalyst-h.

ST transition metal complex **olefin polymn** catalyst; iron complex **ethylene polymn** catalyst; cobalt complex **ethylene polymn** catalyst; bisimidino compd transition metal complex

IT Aluminoxanes

RL: CAT (Catalyst use); USES (Uses)  
 (Me, cocatalyst; transition metal complexes with bisimidino ligands as **olefin polymn.** catalysts)

IT Silica gel, uses

RL: CAT (Catalyst use); USES (Uses)  
 (catalyst support; transition metal complexes with bisimidino ligands as **olefin polymn.** catalysts)

IT Polymerization catalysts

(transition metal complexes with bisimidino ligands as **olefin**

polymn. catalysts)

IT Polyolefins  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (transition metal complexes with bisimidino ligands as **olefin**  
**polymn. catalysts**)

IT 118612-00-3, N,N-Dimethylanilinium tetrakis(pentafluorophenyl)borate  
 RL: CAT (Catalyst use); USES (Uses)  
 (cocatalyst; transition metal complexes with bisimidino ligands as  
**olefin polymn. catalysts**)

---

IT 289708-74-3P 289708-75-4P 289708-76-5P  
289708-77-6P 289708-81-2P 289708-82-3P  
 328239-71-0P 328239-72-1P 328239-73-2P  
 328239-74-3P 328239-75-4P 328239-76-5P  
328239-77-6P 328239-78-7P 328239-79-8P  
 328239-81-2P  
 RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);  
 USES (Uses)  
 (prepn. of transition metal complexes with bisimidino **ligands**  
 )

IT 78-94-4, Methyl vinyl ketone, reactions 110-13-4, 2,5-Hexanedione  
 529-20-4, o-Tolualdehyde 583-05-1, 1-Phenyl-1,4-pentanedione 765-71-9,  
 1-Amino-2,5-dimethylpyrrole 786-98-1 932-95-6, 2,5-  
 Thiophenedicarboxaldehyde 1068-57-1, Acetyl hydrazide 1129-30-2,  
 2,6-Diacetylpyridine 3530-15-2, 4-Amino-3,5-dimethyl-1,2,4-triazole  
 13901-85-4, 6-Methyl-2,5-heptanedione 17223-85-7, N-Aminocarbazole  
 53406-41-0 289708-63-0  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (prepn. of transition metal complexes with bisimidino ligands)

---

IT 3042-24-8P 3042-25-9P 33451-79-5P 195137-30-5P, 1-(2-Methylphenyl)-  
 1,4-pentanedione 289708-72-1P 289708-73-2P 289708-86-7P  
 289709-39-3P 328239-59-4P 328239-60-7P 328239-61-8P 328239-62-9P  
 328239-63-0P 328239-64-1P 328239-65-2P 328239-66-3P 328239-67-4P  
 328239-68-5P 328239-69-6P 328239-70-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (prepn. of transition metal complexes with bisimidino ligands)

---

IT 289708-96-9P 328239-80-1P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of transition metal complexes with bisimidino **ligands**  
 )

IT 9002-88-4P, Polyethylene 9003-07-0P, Polypropylene 25213-02-9P,  
 Ethylene-1-hexene copolymer  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (transition metal complexes with bisimidino ligands as **olefin**  
**polymn. catalysts**)

RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD


RE

(1) Dorer Birgit Angelika; WO 9912981 A 1999 HCPLUS

IT 289708-74-3P 289708-75-4P 289708-76-5P  
 289708-77-6P 289708-81-2P 289708-82-3P  
 328239-71-0P 328239-72-1P 328239-73-2P  
 328239-74-3P 328239-75-4P 328239-76-5P  
328239-77-6P 328239-78-7P 328239-79-8P  
 RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);  
 USES (Uses)  
 (prepn. of transition metal complexes with bisimidino **ligands**  
 )


RN 289708-74-3 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,5-  
 dimethyl-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)




RN 289708-75-4 HCAPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2,5-dimethyl-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)



RN 289708-76-5 HCAPLUS


CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2,5-bis(1-methylethyl)-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)



RN 289708-77-6 HCAPLUS


KATHLEEN FULLER EIC 1700/LAW LIBRARY 308-4290

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2,5-bis(1-methylethyl)-1H-pyrrol-1-amine-.kappa.NN1]]- (9CI) (CA INDEX NAME)




RN 289708-81-2 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2-methyl-5-phenyl-1H-pyrrol-1-amine-.kappa.NN1]]- (9CI) (CA INDEX NAME)




RN 289708-82-3 HCAPLUS


CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2-methyl-5-phenyl-1H-pyrrol-1-amine-.kappa.NN1]]- (9CI) (CA INDEX NAME)




RN 328239-71-0 HCAPLUS  
CN Iron, trichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,5-dimethyl-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)



RN 328239-72-1 HCAPLUS  
CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2-methyl-5-(1-methylethyl)-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)




RN 328239-73-2 HCAPLUS  
CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2-methyl-5-(1-methylethyl)-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)




RN 328239-74-3 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2-methyl-5-(2-methylphenyl)-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)




RN 328239-75-4 HCAPLUS


CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2-methyl-5-(2-methylphenyl)-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)




RN 328239-76-5 HCAPLUS  
CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliidyne]bis[2-methyl-1H-indol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)



RN 328239-77-6 HCAPLUS  
CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliidyne]bis[2-methyl-1H-indol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)




RN 328239-78-7 HCAPLUS  
CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliidyne]bis[9H-carbazol-9-amine-.kappa.NN9]- (9CI) (CA INDEX NAME)



RN 328239-79-8 HCAPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[9H-carbazol-9-amine-.kappa.NN9]- (9CI) (CA INDEX NAME)




IT 289708-96-9P 328239-80-1P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of transition metal complexes with bisimidino ligands  
)


RN 289708-96-9 HCAPLUS

CN Iron, trichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2,5-bis(1-methylethyl)-1H-pyrrol-1-amine-.kappa.NN1]- (9CI) (CA INDEX NAME)



RN 328239-80-1 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[3,5-dimethyl-4H-1,2,4-triazol-4-amine-.kappa.NN4]]- (9CI) (CA INDEX NAME)



L31 ANSWER 7 OF 17 HCAPLUS COPYRIGHT 2002 ACS

AN 2001:115157 HCAPLUS

DN 134:178960

TI Metal complexes comprising a 2,6-diacylpyridine ligand and their use as ethylene polymerization catalysts

IN Sommazzi, Anna; Milani, Barbara; Proto, Antonio; Corso, Gianni; Mestroni, Giovanni; Masi, Francesco

PA Enichem S.p.A., Italy

SO PCT Int. Appl., 58 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07F015-02

ICS C07D213-53; C07F015-06; C08F010-02; C08F004-70

CC 35-3 (Chemistry of Synthetic High Polymers)  
Section cross-reference(s): 78

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001010875	A1	20010215	WO 2000-EP7549	20000803

W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW		
IT 99MI1764	A1 20010205	IT 1999-MI1764	19990805
GB 2369120	A1 20020522	GB 2002-4115	20000803
FI 2002000195	A 20020201	FI 2002-195	20020201
SE 2002000323	A 20020320	SE 2002-323	20020205
PRAI IT 1999-MI1764	A 19990805		
WO 2000-EP7549	W 20000803		
OS MARPAT 134:178960			
AB	An organometallic complex (Q)M(X) <sub>n</sub> (M = transition metal in pos. oxidn. state; Q = 2,6-diacylpyridine monoimine; each X = anionic ligand; n as necessary for neutrality of charge) can be used for the formation of catalytic systems. The complex is prep'd. by relatively simple methods and can be used, combined with a suitable cocatalyst such as an aluminoxane, as a catalyst in (co)polymn. of .alpha.-olefins, esp. ethylene. Thus, 2,6-diacetylpyridine was condensed 1:1 with 2,6-diisopropylaniline to give a monoimine, which was complexed with CoCl <sub>2</sub> . Polymn. of ethylene (0.7 MPa) with this complex and Me aluminoxane (Al/Co = 80) in toluene at 25 .fwdarw. 42.degree. for 30 min gave polyethylene of Mw 12,673 and Mw/Mn 7.5 at 150 g/mmol Co, compared with 11,453, 3.2, and 330 g/mmol Co at 25 .fwdarw. 54.degree. for a catalyst contg. the diimine as ligand.		
ST	ethylene polymn catalyst; diacylpyridine monoimine transition metal complex; polyolefin manuf transition metal complex catalyst		
IT	Aluminoxanes		
	RL: CAT (Catalyst use); USES (Uses)	(Me, cocatalyst, Eurecen 5100-10T; metal complexes comprising a diacylpyridine monoimine ligand as polymn. catalysts)	
IT	Polymerization catalysts		
	(metal complexes comprising a diacylpyridine monoimine ligand as olefin polymn. catalysts)		
IT	Polyolefins		
	RL: IMF (Industrial manufacture); PREP (Preparation)	(metal complexes comprising a diacylpyridine monoimine ligand as polymn. catalysts)	
IT	326613-35-8P 326613-37-0P 326613-40-5P		
	RL: CAT (Catalyst use); IMF (Industrial manufacture); PREP (Preparation); USES (Uses)	(metal complexes comprising a diacylpyridine monoimine ligand as polymn. catalysts)	
IT	9002-88-4P, Polyethylene		
	RL: IMF (Industrial manufacture); PREP (Preparation)	(metal complexes comprising a diacylpyridine monoimine ligand as polymn. catalysts)	
IT	62-53-3, Aniline, reactions		
	RL: RCT (Reactant); RACT (Reactant or reagent)	(prepn. of diacylpyridine imines)	
IT	55137-80-9P, 2,6-Bis[1-(phenylimino)ethyl]pyridine		
	RL: SPN (Synthetic preparation); PREP (Preparation)	(prepn. of diacylpyridine imines)	
IT	263705-60-8P 326613-32-5P		
	RL: IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)	(prepn. of metal complexes comprising a diacylpyridine monoimine ligand as polymn. catalysts)	

IT 88-05-1, 2,4,6-Trimethylaniline 1129-30-2, 2,6-Diacetylpyridine  
 6310-21-0, 2-tert-Butylaniline 24544-04-5, 2,6-Diisopropylaniline  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (prepn. of metal complexes comprising a diacylpyridine monoimine ligand  
 as polymn. catalysts)

IT 326613-30-3P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of metal complexes comprising a diacylpyridine monoimine ligand  
 as polymn. catalysts)


RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Du Pont; WO 9827124 A 1998 HCPLUS  
 (2) Luks; COLLECT CZECH CHEM COMMUN 1998, V63(3), P371 HCPLUS  
 IT 326613-35-8P 326613-37-0P 326613-40-5P  
 RL: CAT (Catalyst use); IMF (Industrial manufacture); PREP (Preparation);  
 USES (Uses)  
 (metal complexes comprising a diacylpyridine monoimine ligand  
 as polymn. catalysts)


RN 326613-35-8 HCPLUS

CN Cobalt, [1-[6-[1-[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]ethyl]-2-  
 pyridinyl-.kappa.N]ethanone-.kappa.O]dichloro- (9CI) (CA INDEX NAME)




RN 326613-37-0 HCPLUS

CN Cobalt, dichloro[1-[6-[1-[(2,4,6-trimethylphenyl)imino-.kappa.N]ethyl]-2-  
 pyridinyl-.kappa.N]ethanone-.kappa.O]- (9CI) (CA INDEX NAME)



RN 326613-40-5 HCPLUS

CN Iron, [1-[6-[1-[2,6-bis(1-methylethyl)phenyl]imino-.kappa.N]ethyl]-2-  
 pyridinyl-.kappa.N]ethanone-.kappa.O]dichloro- (9CI) (CA INDEX NAME)



L31 ANSWER 8 OF 17 HCAPLUS COPYRIGHT 2002 ACS

AN 2001:78431 HCAPLUS

DN 134:148001

TI Transition metal complexes with tridentate ligands and their use as catalysts in manufacture of olefin (co)polymers

IN Luinstra, Gerrit; Werne, Gerald; Rief, Ursula; Kristen, Marc Oliver; Queisser, Joachim; Geprags, Michael

PA BASF Aktiengesellschaft, Germany

SO PCT Int. Appl., 31 pp.

CODEN: PIXXD2

DT Patent

LA German

IC ICM C08F010-00

ICS C08F004-70; C08F210-00; C07F015-02; C07F015-06; C07F015-00

CC 35-3 (Chemistry of Synthetic High Polymers)


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001007491	A1	20010201	WO 2000-EP6559	20000711
	W: JP, KR, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				

PRAI DE 1999-19934463 A 19990727

OS MARPAT 134:148001


GI




AB The complexes have the formula I [A is a noncoordinating or barely coordinating anion; M = Fe, Ru, Co, Rh; Q, T = neutral or monoanionic monodentate ligand; R1, R2 = C6-16 aryl with halo, nitro, cyano, sulfo or trihalomethyl substituents in both positions ortho to the bond to Na or

Nb; R3, R4 = H, C1-10 alkyl, C3-10 cycloalkyl, C6-16 aryl, alkylaryl with 1-10 C atoms in the alkyl portion and 6-14 C atoms in the aryl portion, SiR83; R5-R7 = H, C1-10 alkyl, C3-10 cycloalkyl, C6-16 aryl, alkylaryl with 1-10 C atoms in the alkyl portion and 6-14 C atoms in the aryl portion, SiR83, functional groups consisting of Group IVA-VIIA elements, or R5R6 and/or R6R7 form a five- to seven-membered (un)substituted fused ring; R8 = C1-10 alkyl, C3-10 cycloalkyl, C6-16 aryl, alkylaryl with 1-10 C atoms in the alkyl portion and 6-14 C atoms in the aryl portion; m, p = 0-3; n, q = 1-3]. .alpha.-Olefins are homo- or copolymerd. in the presence of .gtoreq.1 I and a cocatalyst consisting of a strong neutral Lewis acid or an ionic compd. having a Lewis or Bronsted-acid-cation. Thus, 2,6-dichloroaniline was condensed 2:1 with 2,6-diacetylpyridine and the product treated with FeCl<sub>2</sub> to give a I, which was used with Me aluminoxane to polymerize ethylene to a homopolymer with Mw/Mn 3.01.

ST transition metal complex olefin polymn catalyst  
 IT Aluminoxanes  
 RL: CAT (Catalyst use); USES (Uses)  
 (Me, cocatalyst; transition metal complexes with tridentate ligands as olefin polymn. catalysts)  
 IT Lewis acids  
 RL: CAT (Catalyst use); USES (Uses)  
 (cocatalysts; transition metal complexes with tridentate ligands as olefin polymn. catalysts)  
 IT Polymerization catalysts  
 (transition metal complexes with tridentate ligands as olefin polymn. catalysts)  
 IT 323179-69-7P 323179-71-1P  
 RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation); USES (Uses)  
 (prepn. of transition metal complexes with tridentate ligands as olefin polymn. catalysts)  
 IT 608-31-1, 2,6-Dichloroaniline 1129-30-2, 2,6-Diacetylpyridine  
 6968-24-7, 2,6-Dibromo-4-methylaniline  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (prepn. of transition metal complexes with tridentate ligands as olefin polymn. catalysts)  
 IT 323179-67-5P 323179-70-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. of transition metal complexes with tridentate ligands as olefin polymn. catalysts)  
 IT 9002-88-4P, Polyethylene 25213-02-9P, Ethylene-1-hexene copolymer  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (transition metal complexes with tridentate ligands as olefin polymn. catalysts)  
 RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE  
 (1) Brookhart, M; WO 9847934 A 1998 HCPLUS  
 (2) Dorer, B; WO 9912981 A 1999 HCPLUS  
 (3) Du Pont; WO 9827124 A 1998 HCPLUS  
 IT 323179-69-7P 323179-71-1P  
 RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation); USES (Uses)  
 (prepn. of transition metal complexes with tridentate ligands as olefin polymn. catalysts)  
 RN 323179-69-7 HCPLUS  
 CN Iron, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-dibromo-4-methylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)



RN 323179-71-1 HCPLUS  
 CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-dichlorobenzene-.kappa.N]]- (9CI) (CA INDEX NAME)



L31 ANSWER 9 OF 17 HCPLUS COPYRIGHT 2002 ACS  
 AN 2000:909415 HCPLUS  
 DN 134:208205  
 TI **Ethylene polymerization** and copolymerization with higher .alpha.-olefins using catalysts based on iron chlorides containing 2,6-bis(imino)pyridine ligands  
 AU Ivancheva, N. I.; Badaev, V. K.; Oleinik, I. I.; Ivanchev, S. S.; Tolstikov, G. A.  
 CS S.-Peterb. Fil. Inst. Kataliza im. G. K. Boreskova, Sib. Otd. Ross. Akad. Nauk, St. Petersburg, Russia  
 SO Doklady Akademii Nauk (2000), 374(5), 648-650  
 CODEN: DAKNEQ; ISSN: 0869-5652  
 PB MAIK Nauka  
 DT Journal  
 LA Russian  
 CC 35-3 (Chemistry of Synthetic High Polymers)  
 AB Four new 2,6-bis[(4- or/and 6-alkyl-2-cyclopentylphenylimino)pyridine] iron(II) dichloride complexes (alkyl - H or Me) were tested as catalysts for **ethylene polymn.** and copolymn. with 1-octene and

4-methylpentene. The catalysts yielded polyethylene with yields comparable to those obtained on the metallocene catalysts.

ST iron bisiminopyridine chloride catalyst **ethylene polymn**  
octene methylpentene

IT **Polymerization catalysts**  
Substituent effects

(**ethylene polymn.** and copolymn. with .alpha.-  
olefins using FeCl<sub>2</sub>-based catalysts contg. 2,6-  
bis(imino)pyridine ligands)

IT 328313-03-7 328313-05-9 328313-06-0  
328313-08-2

RL: CAT (Catalyst use); USES (Uses)

(**ethylene polymn.** and copolymn. with .alpha.-  
olefins using FeCl<sub>2</sub>-based catalysts contg. 2,6-  
bis(imino)pyridine ligands)

IT 9002-88-4P, Polyethylene 26221-73-8P, **Ethylene-1-octene**  
copolymer 130315-97-8P, **Ethylene-4-methylpentene copolymer**

RL: SPN (Synthetic preparation); PREP (Preparation)  
(**ethylene polymn.** and copolymn. with .alpha.-  
olefins using FeCl<sub>2</sub>-based catalysts contg. 2,6-  
bis(imino)pyridine ligands)


IT 328313-03-7 328313-05-9 328313-06-0  
328313-08-2

RL: CAT (Catalyst use); USES (Uses)

(**ethylene polymn.** and copolymn. with .alpha.-  
olefins using FeCl<sub>2</sub>-based catalysts contg. 2,6-  
bis(imino)pyridine ligands)



RN 328313-03-7 HCPLUS

CN Iron, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2-  
cyclopentylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)




RN 328313-05-9 HCPLUS

CN Iron, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2-  
cyclopentyl-6-methylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)



RN 328313-06-0 HCAPLUS


CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2-cyclopentyl-4,6-dimethylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)




RN 328313-08-2 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2,6-dicyclopentylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



L31 ANSWER 10 OF 17 HCPLUS COPYRIGHT 2002 ACS  
 AN 2000:785358 HCPLUS  
 DN 134:86580  
 TI The behavior of homogeneous iron-based catalysts bearing pyridine diimine ligands for **ethylene polymerization**  
 AU Qiu, Jiao-Ming; Sun, Ling-Gang; Hu, You-Liang; Li, Yu-Fei  
 CS State Key Laboratory of Engineering Plastics, Center for Molecular Science Institute of Chemistry, Chinese Academy of Sciences, Beijing, 100080, Peop. Rep. China  
 SO Chinese Journal of Polymer Science (2000), 18(6), 509-513  
 CODEN: CJPSEG; ISSN: 0256-7679  
 PB Springer-Verlag  
 DT Journal  
 LA English  
 CC 35-3 (Chemistry of Synthetic High Polymers)  
 Section cross-reference(s): 28, 78  
 AB The **polymn.** of **ethylene** by two iron-based catalysts, [2,6-ArN: CMe<sub>2</sub>C<sub>5</sub>H<sub>3</sub>N]FeCl<sub>2</sub> with Ar = 2,6-Me<sub>2</sub>C<sub>6</sub>H<sub>3</sub> (I) and with Ar = 2,6-iso-Pr<sub>2</sub>C<sub>6</sub>H<sub>3</sub> (II) has been investigated. II in conjunction with Me aluminoxane produces higher mol. wt. polyethylene (PE) and broadened polydispersities relative to I under analogous conditions and all polymers are linear. The kinetic profiles with both catalysts showed a smooth pattern during both rate build-up and rate lowering, which are different from metallocene catalysts. The polymn. activity increases with Al/Fe ratio and an optimum temp. range of 40.apprx.45.degree.C was obsd. The PE mol. wt. decreases with increasing Al/Fe ratio and polymn. temp.  
 ST iron complex pyridine diimine catalyst polymn; **ethylene**  
**polymn** iron complex catalyst

IT Aluminoxanes  
RL: CAT (Catalyst use); USES (Uses)  
(Me, cocatalysts; homogeneous iron-based catalysts bearing pyridine diimine ligands for polymn. of ethylene)

IT Polymerization kinetics  
(of ethylene in presence of homogeneous iron-based catalysts bearing pyridine diimine ligands)

IT Polymerization catalysts  
(prepn. of homogeneous iron-based catalysts bearing pyridine diimine ligands for polymn. of ethylene)

IT 74-85-1, Ethylene, reactions  
RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)  
(homogeneous iron-based catalysts bearing pyridine diimine ligands for polymn. of)

IT 9002-88-4P, Polyethylene  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(homogeneous iron-based catalysts bearing pyridine diimine ligands for prepn. of)

IT 204203-14-5P 204203-16-7P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(intermediate; prepn. of homogeneous iron-based catalysts bearing pyridine diimine ligands for polymn. of ethylene)

IT 204203-10-1P 207129-93-9P  
RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);  
USES (Uses)  
(prepn. of homogeneous iron-based catalysts bearing pyridine-diimine ligands for polymn. of ethylene)

IT 87-62-7, 2,6-Dimethylaniline 7758-94-3, Ferrous chloride 20857-17-4,  
2,4-Diacetylpyridine 24544-04-5, 2,6-Diisopropylaniline  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(starting material; prepn. of homogeneous iron-based catalysts bearing pyridine diimine ligands for polymn. of ethylene)

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD


RE

- (1) Britovsek, G; Chem Commun 1998, P849
- (2) Brookhart, M; WO 9830612 1998 HCPLUS
- (3) Chien, J; J Polym Sci Part A 1986, V24, P2483 HCPLUS
- (4) Chien, J; J Polym Sci Part A 1988, V26, P2369 HCPLUS
- (5) Freenatle, M; C & EN 1998, P11
- (6) Johnson, L; J Am Chem Soc 1996, V118, P267 HCPLUS
- (7) Killian, C; J Am Chem Soc 1995, V117, P6414
- (8) Killian, C; J Am Chem Soc 1996, V118, P11664 HCPLUS
- (9) Reddy, S; Prog Polym Sci 1995, V20, P309 HCPLUS
- (10) Small, B; J Am Chem Soc 1998, V120, P4049 HCPLUS
- (11) Small, B; Macromolecules 1999, V32, P2120 HCPLUS

IT 204203-10-1P 207129-93-9P  
RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);  
USES (Uses)  
(prepn. of homogeneous iron-based catalysts bearing pyridine diimine ligands for polymn. of ethylene)


RN 204203-10-1 HCPLUS

CN Iron, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)



RN 207129-93-9 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2,6-dimethylbenzene-.kappa.N]], (SP-5-13)- (9CI) (CA INDEX NAME)



L31 ANSWER 11 OF 17 HCAPLUS COPYRIGHT 2002 ACS

AN 2000:384252 HCAPLUS

DN 133:5119

TI Catalysts for olefin polymerization and process for producing olefin polymers

IN Okuda, Fumio

PA Idemitsu Petrochemical Co., Ltd., Japan

SO PCT Int. Appl., 27 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

IC ICM C08F004-70

ICS C08F010-00

CC 35-3 (Chemistry of Synthetic High Polymers)  
Section cross-reference(s): 67

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000032643	A1	20000608	WO 1999-JP6768	19991202
	W: JP, KR, US				

RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE

EP 1054022 A1 20001122 EP 1999-973041 19991202

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI

PRAI JP 1998-342458 A 19981202

WO 1999-JP6768 W 19991202

OS MARPAT 133:5119

AB The catalysts comprise a compd. of a transition metal in Groups 8-10 of the Periodic Table having a nitrogenous tridentate ligand and an organoaluminum compd. such as trimethylaluminum. The catalysts are highly active and enable a high-mol. polyolefin to be produced. Consequently, a high-mol. polyolefin (esp. polyethylene) suitable for practical use can be industrially advantageously produced without using methylaluminoxane, which is expensive, has poor handleability and poor storage stability, and is highly dangerous.

ST ethylene polymn catalyst; nitrogen tridentate ligand transition metal complex catalyst; polyolefin high mol wt prepn

IT Polymerization catalysts  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for olefin polymn.)

IT Transition metal complexes  
RL: CAT (Catalyst use); USES (Uses)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for olefin polymn.)

IT Polyolefins  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for olefin polymn.)

IT 75-24-1, Trimethylaluminum 100-99-2, Triisobutylaluminum, uses  
**204203-10-1**  
RL: CAT (Catalyst use); USES (Uses)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for olefin polymn.)

IT 9002-88-4P, Polyethylene  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for olefin polymn.)

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD


RE

- (1) E I Du Pont de Nemours And Company; US 5955555 A HCPLUS
- (2) E I Du Pont de Nemours And Company; EP 946609 A1 HCPLUS
- (3) E I Du Pont de Nemours And Company; EP 951489 A1 HCPLUS
- (4) E I Du Pont de Nemours And Company; WO 9827124 A 1998 HCPLUS
- (5) E I Du Pont de Nemours And Company; WO 9830612 A 1998 HCPLUS
- (6) Mitsui Chemicals Ltd; JP 107712 A 1998
- (7) Mitsui Chemicals Ltd; JP 20001512 A 2000
- (8) Mitsui Chemicals Ltd; JP 20001513 A 2000

IT **204203-10-1**  
RL: CAT (Catalyst use); USES (Uses)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for olefin polymn.)

RN 204203-10-1 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]-, (SP-5-13)- (9CI) (CA INDEX NAME)



L31 ANSWER 12 OF 17 HCPLUS COPYRIGHT 2002 ACS

AN 2000:384251 HCPLUS

DN 133:17992

TI Catalysts for **olefin polymerization** and process for producing **olefin** polymers

IN Okuda, Fumio; Sato, Haruhito; Kuramoto, Masahiko

PA Idemitsu Petrochemical Co., Ltd., Japan

SO PCT Int. Appl., 52 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

IC ICM C08F004-70

ICS C08F010-00

CC 35-3 (Chemistry of Synthetic High Polymers)

Section cross-reference(s): 67

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000032642	A1	20000608	WO 1999-JP6767	19991202
	W: JP, KR, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	EP 1054021	A1	20001122	EP 1999-957408	19991202
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				

PRAI JP 1998-342457 A 19981202  
 JP 1998-342459 A 19981202  
 WO 1999-JP6767 W 19991202

OS MARPAT 133:17992

AB The catalysts comprise a compd. of a transition metal in Groups 8-10 of the Periodic Table having a nitrogenous tridentate ligand, a clay, clay mineral, or lamellar ion-exchanging compd., an organosilane compd., an organoaluminum compd., etc. The catalysts are highly active, do not adhere to reactor walls, and can give a polyolefin excellent in powder morphol. Consequently, a polyolefin (esp. polyethylene) can be industrially advantageously produced.

ST **ethylene polymn** catalyst; nitrogen tridentate ligand  
 transition metal complex catalyst; polyolefin high mol wt prepn

IT Polymerization catalysts  
 (highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for **olefin polymn.**)

IT Clays, uses  
Transition metal complexes  
RL: CAT (Catalyst use); USES (Uses)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for **olefin polymn.**)

IT Polyolefins  
RL: IMF (Industrial manufacture); PRP (Properties); PREP (Preparation)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for **olefin polymn.**)

IT Ion exchangers  
(lamellar; highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for **olefin polymn.**)

IT 75-24-1, Trimethylaluminum 100-99-2, Triisobutylaluminum, uses 1318-93-0, Montmorillonite, uses 7786-30-3, Magnesium chloride, uses 114502-16-8 187247-40-1, Kunipia F 204203-10-1  
RL: CAT (Catalyst use); USES (Uses)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for **olefin polymn.**)

IT 9002-88-4P, Polyethylene  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for **olefin polymn.**)

RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE


- (1) E I Du Pont de Nemours And Company; US 5955555 A HCAPLUS
- (2) E I Du Pont de Nemours And Company; EP 946609 A1 HCAPLUS
- (3) E I Du Pont de Nemours And Company; EP 951489 A1 HCAPLUS
- (4) E I Du Pont de Nemours And Company; WO 9827124 A 1998 HCAPLUS
- (5) E I Du Pont de Nemours And Company; WO 9830612 A 1998 HCAPLUS
- (6) Idemitsu Petrochem Co Ltd; JP 11269222 A HCAPLUS
- (7) Idemitsu Petrochem Co Ltd; JP 11269223 A HCAPLUS
- (8) Idemitsu Petrochem Co Ltd; JP 11269224 A HCAPLUS
- (9) Idemitsu Petrochem Co Ltd; WO 9948930 A 1999 HCAPLUS
- (10) Mitsubishi Kasei Corporation; JP 05105721 A HCAPLUS
- (11) Mitsubishi Kasei Corporation; JP 05301917 A HCAPLUS
- (12) Mitsubishi Kasei Corporation; US 5308811 A HCAPLUS
- (13) Mitsubishi Kasei Corporation; EP 698621 A1 HCAPLUS
- (14) Mitsubishi Kasei Corporation; EP 511665 A2 1992 HCAPLUS
- (15) Mitsui Chemicals Ltd; JP 20001513 A 2000

IT 204203-10-1

RL: CAT (Catalyst use); USES (Uses)  
(highly active transition metal complex catalysts contg. nitrogenous tridentate ligands for **olefin polymn.**)

RN 204203-10-1 HCAPLUS

CN Iron, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)



L31 ANSWER 13 OF 17 HCAPLUS COPYRIGHT 2002 ACS  
 AN 1999:577442 HCAPLUS  
 DN 131:337391  
 TI Iron and Cobalt **Ethylene Polymerization** Catalysts  
 Bearing 2,6-Bis(Imino)Pyridyl Ligands: Synthesis, Structures, and  
 Polymerization Studies  
 AU Britovsek, George J. P.; Bruce, Michael; Gibson, Vernon C.; Kimberley,  
 Brian S.; Maddox, Peter J.; Mastroianni, Sergio; McTavish, Stuart J.;  
 Redshaw, Carl; Solan, Gregory A.; Stroemberg, Staffan; White, Andrew J.  
 P.; Williams, David J.  
 CS Department of Chemistry, Imperial College, South Kensington London, SW7  
 2AY, UK  
 SO Journal of the American Chemical Society (1999), 121(38), 8728-8740  
 CODEN: JACSAT; ISSN: 0002-7863  
 PB American Chemical Society  
 DT Journal  
 LA English  
 CC 35-3 (Chemistry of Synthetic High Polymers)  
 Section cross-reference(s): 75, 78  
 AB The synthesis, characterization, and **ethylene polymn.**  
 behavior of a series of iron and cobalt halide complexes,  $LMX_n$  ( $M = Fe, X = Cl, n = 2, 3, X = Br, n = 2; M = Co, X = Cl, n = 2$ ), bearing chelating 2,6-bis(imino)pyridyl ligands  $L$  [ $L = 2,6-(ArNCR1)2C5H3N$ ] is reported.  
 X-ray diffraction studies show the geometry at the metal centers to be either distorted square pyramidal or distorted trigonal-bipyramidal.  
 Treatment of the complexes  $LMX_n$  with methylaluminoxane (MAO) leads to highly active **ethylene polymn.** catalysts converting **ethylene** to highly linear polyethylene (PE).  $LFEx2$  precatalysts with ketimine ligands ( $R1 = Me$ ) are approx. an order of magnitude more active than precatalysts with aldimine ligands ( $R1 = H$ ). Catalyst productivities in the range 3,750-20,600 g/mmol.cntdot.h.cntdot.bar are obsd. for Fe-based ketimine catalysts, while Co ketimine systems display activities of 450-1740 g/mmol.cntdot.h.cntdot.bar. Mol. wts. ( $M_w$ ) of the polymers produced are in the range 14,000-611,000. Changing reaction conditions also affects productivity and mol. wt.; in some systems, a bimodal mol. wt. distribution is obsd. On the basis of evidence gathered to date, the lower mol. wt. fraction is a result of chain transfer to aluminum while the higher mol. wt. fraction is produced by a combination of mainly  $\beta$ -H transfer and some chain transfer to aluminum.  
 ST iron cobalt **ethylene polymn** catalyst; iminopyridyl  
 contg iron cobalt polymn catalyst; polyethylene prepn iminopyridyl iron

cobalt catalyst; methylaluminoxane **ethylene polymn**  
 catalyst iron cobalt

IT Aluminoxanes  
 RL: CAT (Catalyst use); USES (Uses)  
 (Me, cocatalysts; prepn. of iron and cobalt **ethylene polymn**. catalysts bearing bis(imino)pyridyl ligands for **polymn. of ethylene**)

IT Bond angle  
 Bond length  
 Crystal structure  
Molecular structure  
 (of iron and cobalt catalysts bearing bis(imino)pyridyl ligands for **polymn. of ethylene**)

IT Polymerization catalysts  
 (prepn. of iron and cobalt **ethylene polymn**. catalysts bearing bis(imino)pyridyl ligands for **polymn. of ethylene**)

IT 204203-10-1P 207129-94-0P 207129-97-3P  
 247911-89-3P  
 RL: CAT (Catalyst use); PRP (Properties); SPN (Synthetic preparation);  
 PREP (Preparation); USES (Uses)  
 (catalysts; prepn. of iron and cobalt **ethylene polymn**. catalysts bearing bis(imino)pyridyl ligands for **polymn. of ethylene**)

IT 204203-12-3P 207129-93-9P 207129-96-2P  
 210155-45-6P 210768-87-9P 221391-12-4P  
 221391-13-5P 221391-16-8P 247911-91-7P  
 247911-92-8P  
 RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);  
 USES (Uses)  
 (catalysts; prepn. of iron and cobalt **ethylene polymn**. catalysts bearing bis(imino)pyridyl ligands for **polymn. of ethylene**)

IT 204203-14-5P 204203-16-7P 204203-17-8P 210155-39-8P 210155-42-3P  
 219729-70-1P 221391-10-2P 221391-11-3P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and characterization of bis(imino)pyridyl ligands for iron and cobalt **polymn. catalysts for ethylene**)

IT 9002-88-4P, Polyethylene  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of iron and cobalt **ethylene polymn**. catalysts bearing bis(imino)pyridyl ligands for **polymn. of ethylene**)

IT 7646-79-9, Cobalt dichloride, reactions 7758-94-3, Iron dichloride  
 7789-46-0, Iron dibromide  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reaction with bis(imino)pyridyl ligands)

IT 1129-30-2, 2,6-Diacetylpyridine 24544-04-5, 2,6-Diisopropylaniline  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reaction with diisopropylaniline)

RE.CNT 70 THERE ARE 70 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Alcock, N; J Chem Soc 1984, P1937 HCPLUS
- (2) Atiqullah, M; Eur Polym J 1998, V34, P1511 HCPLUS
- (3) Bennett, A; 1998 HCPLUS
- (4) Bennett, A; WO 9827124 1998 HCPLUS
- (5) Bennett, A; CHEMTECH 1999
- (6) Blandamer, M; J Chem Soc 1978, P1001 HCPLUS
- (7) Bochmann, M; Angew Chem Int Ed 1994, V33, P1634

(8) Bochmann, M; J Chem Soc 1996, P255 HCPLUS  
(9) Brintzinger, H; Angew Chem Int Ed 1995, V34, P1143 HCPLUS  
(10) Britovsek, G; 1999 HCPLUS  
(11) Britovsek, G; WO 9912981 1999 HCPLUS  
(12) Britovsek, G; Angew Chem Int Ed 1999, V38, P428 HCPLUS  
(13) Britovsek, G; Chem Commun 1998, P849  
(14) Brookhart, M; HCPLUS  
(15) Brookhart, M; WO 9830612 1998 HCPLUS  
(16) Bruce, M; Chem Commun 1998, P2523 HCPLUS  
(17) Busico, V; Macromolecules 1994, V27, P7538 HCPLUS  
(18) Cetinkaya, B; J Mol Catal A: Chem 1999, V142, P101 HCPLUS

---


(19) Chien, J; J Polym Sci A 1990, V28, P15 HCPLUS  
(20) Cossee, P; J Catal 1964, V3, P80 HCPLUS  
(21) Curry, J; Inorg Chem 1967, V6, P1570 HCPLUS  
(22) Edwards, D; J Chem Soc 1990, P3161 HCPLUS  
(23) Edwards, D; Polyhedron 1992, V11, P1569 HCPLUS  
(24) Evans, D; J Chem Soc 1959, P2003 HCPLUS  
(25) Evans, D; J Chem Soc 1988, P2927 HCPLUS  
(26) Figgins, P; J Am Chem Soc 1960, V82, P820 HCPLUS  
(27) Fink, G; Ziegler Catalysts: Recent Scientific Innovations and Technological Improvement 1995  
(28) Fogg, P; Solubility of gases in liquids 1991  
(29) Gibson, V; Unpublished results  
(30) Haarman, H; Organometallics 1997, V16, P54 HCPLUS  
(31) Haarman, H; Organometallics 1997, V16, P887 HCPLUS  
(32) Haarman, H; Organometallics 1997, V16, P979 HCPLUS  
(33) Haslam, J; Identification and Analysis of Plastic 2nd ed 1972  
(34) Havan, R; Chem Ind 1997, P212  
(35) Heinemann, J; Macromol Chem Phys 1999, V200, P384 HCPLUS  
(36) Johnson, L; 1996 HCPLUS  
(37) Johnson, L; WO 9623010 1996 HCPLUS  
(38) Johnson, L; J Am Chem Soc 1995, V117, P6414 HCPLUS  
(39) Johnson, L; J Am Chem Soc 1996, V118, P267 HCPLUS  
(40) Jordan, R; Adv Organomet Chem 1991, V32, P325 HCPLUS  
(41) Kaminsky, W; Macromol Chem Phys 1996, V197, P3907 HCPLUS  
(42) Killian, C; J Am Chem Soc 1996, V118, P11664 HCPLUS  
(43) Krumholz, P; Inorg Chem 1965, V4, P612 HCPLUS  
(44) Lions, F; J Am Chem Soc 1957, V79, P2733 HCPLUS  
(45) Marques, M; J Polym Sci A 1993, V31, P219 HCPLUS  
(46) Mecking, S; J Am Chem Soc 1998, V120, P888 HCPLUS  
(47) Morassi, R; Coord Chem Rev 1973, P343 HCPLUS  
(48) Naga, N; Polymer 1998, V39, P5059 HCPLUS  
(49) Onggo, D; Aust J Chem 1991, V44, P331 HCPLUS  
(50) Pellecchia, C; Macromol Rapid Commun 1998, V19, P651 HCPLUS  
(51) Pelletier, J; Angew Chem Int Ed 1996, V35, P1854 HCPLUS  
(52) Pelletier, J; Stud Surf Sci Catal 1994, V89, P249 HCPLUS  
(53) Roberts, G; NMR of Macromolecules: A Practical Approach 1993  
(54) Scheer, C; Polyhedron 1996, V15, P219 HCPLUS  
(55) Schleis, T; J Organomet Chem 1998, V569, P159 HCPLUS  
(56) Siedle, A; Organometallics 1991, V10, P400 HCPLUS  
(57) Siemens Analytical X-ray Instruments Inc; SHELXTL PC version 5.03 1984  
(58) Sinclair, K; Chem Ind 1994, P857 HCPLUS  
(59) Small, B; Am Chem Soc Div Polym Chem 1998, V39, P213 HCPLUS  
(60) Small, B; J Am Chem Soc 1998, V120, P4049 HCPLUS  
(61) Small, B; J Am Chem Soc 1998, V120, P7143 HCPLUS  
(62) Small, B; Macromolecules 1999, V32, P2120 HCPLUS  
(63) Soga, K; Makromol Chem 1993, V194, P1745 HCPLUS  
(64) Stehling, U; Organometallics 1994, V13, P964 HCPLUS  
(65) Stoufer, R; J Am Chem Soc 1956, V78, P6016 HCPLUS  
(66) Suhm, J; Macromol Symp 1998, V129, P1 HCPLUS

(67) Thayer, A; Chem Eng News 1995, P15  
 (68) Tsutsui, T; Polymer 1989, V30, P428 HCPLUS  
 (69) Yamamoto, A; J Am Chem Soc 1968, V90, P1878 HCPLUS  
 (70) Zambelli, A; Macromolecules 1989, V22, P2186 HCPLUS  
 IT 204203-10-1P 207129-94-0P 207129-97-3P  
 247911-89-3P

RL: CAT (Catalyst use); PRP (Properties); SPN (Synthetic preparation);  
 PREP (Preparation); USES (Uses)  
 (catalysts; prepn. of iron and cobalt **ethylene polymn**  
 . catalysts bearing bis(imino)pyridyl ligands for  
**polymn. of ethylene**)


RN 204203-10-1 HCPLUS

CN Iron, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethyliyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)




RN 207129-94-0 HCPLUS

CN Iron, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethyliyne]bis[2,4,6-trimethylbenzenamine-.kappa.N]]-, (TB-5-22)- (9CI) (CA INDEX NAME)




RN 207129-97-3 HCPLUS

CN Cobalt, dichloro[N,N'-[(2,6-pyridinediyl-.kappa.N)diethyliyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)



RN 247911-89-3 HCPLUS

CN Iron, dibromo[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2,4,6-trimethylbenzenamine-.kappa.N]]-, (TB-5-22)- (9CI) (CA INDEX NAME)



IT 204203-12-3P 207129-93-9P 207129-96-2P

210155-45-6P 210768-87-9P 221391-12-4P

221391-13-5P 221391-16-8P 247911-91-7P

247911-92-8P


RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);  
USES (Uses)

(catalysts; prepn. of iron and cobalt ethylene polymn

. catalysts bearing bis(imino)pyridyl ligands for  
polymn. of ethylene)


RN 204203-12-3 HCPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2-(1,1-dimethylethyl)benzenamine]]-, (SP-5-13)- (9CI) (CA INDEX NAME)




RN 207129-93-9 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,6-dimethylbenzylamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)




RN 207129-96-2 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethylidyne]bis[2,6-dimethylbenzylamine-.kappa.N]]- (9CI) (CA INDEX NAME)




RN 210155-45-6 HCAPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,4,6-trimethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)




RN 210768-87-9 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2-(1,1-dimethylethyl)benzenamine-.kappa.N]-, (SP-5-13)- (9CI) (CA INDEX NAME)




RN 221391-12-4 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethylidyne]bis[2,6-diethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)




RN 221391-13-5 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethyliyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)



RN 221391-16-8 HCAPLUS


CN Iron, trichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethyliyne]bis[2,4,6-trimethylbenzenamine-.kappa.N]]-, (OC-6-31)- (9CI) (CA INDEX NAME)



RN 247911-91-7 HCAPLUS


CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethyliyne]bis[2,4,6-

trimethylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)



RN 247911-92-8 HCAPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethylidyne]bis[2,4,6-trimethylbenzenamine-.kappa.N]]- (9CI) (CA INDEX NAME)



L31 ANSWER 14 OF 17 HCAPLUS COPYRIGHT 2002 ACS

AN 1999:428714 HCAPLUS

DN 131:214558

TI Novel, highly active iron and cobalt catalysts for **olefin polymerization**

AU Bennett, Alison M. A.

CS Central Research and Development, E. I. DuPont de Nemours and Co. Inc., USA

SO CHEMTECH (1999), 29(7), 24-28

CODEN: CHTEDD; ISSN: 0009-2703

PB American Chemical Society

DT Journal; General Review

LA English

CC 35-0 (Chemistry of Synthetic High Polymers)

Section cross-reference(s): 67, 78

AB A review, with 36 refs., on iron and cobalt complexes with bulky tridentate ligands, i.e. pyridine bis-imine ligands, activated with MAO, as

chain transfer polymn. catalysts for manuf. of linear polyethylene. Synthesis and structure of the metal complexes, effect of catalyst structure on polymer properties, and flexibility of the catalyst systems are discussed. Prospects of the catalyst systems for alpha.-olefin prodn. are outlined.

ST review iron complex bulky tridentate ligand polymn catalyst; cobalt pyridinebisimine polymn catalyst chain walk review; chain transfer polymn catalyst bulky ligand complex review

IT Aluminoxanes

RL: CAT (Catalyst use); USES (Uses)  
~~(Me; highly active iron and cobalt catalysts with bulky pyridinebisimine ligands for olefin polymn.)~~

IT Polymerization catalysts  
~~(coordination; highly active iron and cobalt catalysts with bulky pyridinebisimine ligands for olefin polymn.)~~

IT Chain transfer  
~~(highly active iron and cobalt catalysts with bulky pyridinebisimine ligands for olefin polymn.)~~

IT 204203-10-1, 2,6-Bis[1-(2,6-diisopropylphenylimino)ethyl]pyridineiron(II) chloride 207129-97-3  
 RL: CAT (Catalyst use); USES (Uses)  
~~(highly active iron and cobalt catalysts with bulky pyridinebisimine ligands for olefin polymn.)~~

IT 9002-88-4P, Polyethylene  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
~~(highly active iron and cobalt catalysts with bulky pyridinebisimine ligands for olefin polymn.)~~

RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Bennett, A; WO 98/27124 1998 HCPLUS
- (2) Brintzinger, H; Angew Chem Int Ed Engl 1995, V34, P1143 HCPLUS
- (3) Britovsek, G; Chem Commun 1998, P849
- (4) Brookhart, M; WO 96/23010 1996 HCPLUS
- (5) Brookhart, M; J Am Chem Soc 1990, V112, P5634 HCPLUS
- (6) Brookhart, M; Macromolecules 1995, V28, P5378 HCPLUS
- (7) Covezzi, M; Macromol Symp 1995, V89, P577 HCPLUS
- (8) Desjardins, S; J Organomet Chem 1996, V515, P233 HCPLUS
- (9) Desjardins, S; J Organomet Chem 1997, V554, P163
- (10) Flory, P; J Am Chem Soc 1940, V62, P1561 HCPLUS
- (11) Johnson, L; J Am Chem Soc 1995, V117, P6414 HCPLUS
- (12) Johnson, L; J Am Chem Soc 1996, V118, P267 HCPLUS
- (13) Kaminsky, W; Macromol Chem Phys 1996, V197, P3907 HCPLUS
- (14) Keim, W; Angew Chem Int Ed Engl 1978, V17, P466
- (15) Keim, W; Angew Chem Int Ed Engl 1981, V20, P116
- (16) Keim, W; Angew Chem Int Ed Engl 1987, V26, P1012
- (17) Lippert, F; WO 96/37522 1996 HCPLUS
- (18) Lippert, F; WO 96/37523 1996 HCPLUS
- (19) Mecking, S; J Am Chem Soc 1998, V120, P888 HCPLUS
- (20) Mohring, P; Organomet Chem 1994, V479, P1
- (21) Mohring, V; Angew Chem Int Ed Engl 1985, V24, P1001
- (22) Natta, G; Angew Chem 1956, V68, P393 HCPLUS
- (23) Schmidt, G; J Am Chem Soc 1985, V107, P1443 HCPLUS
- (24) Schubbe, R; Makromol Phys 1995, V196, P467 HCPLUS
- (25) Schulz, G; Z Phys Chem Abt B 1935, V30, P379
- (26) Schulz, G; Z Phys Chem Abt B 1939, V43, P25
- (27) Small, B; J Am Chem Soc 1998, V120, P4049 HCPLUS
- (28) Small, B; J Am Chem Soc 1998, V120, P7143 HCPLUS
- (29) Small, B; Polym Prepr Am Chem Soc Div Polym Chem 1998, V39, P213 HCPLUS
- (30) Small, B; to be published in Macromolecules
- (31) Starzewski, K; Transition Metals and Organometallics as Catalysts for


Olefin Polymerization 1988, P349 HCAPLUS  
 (32) Timonen, S; J Mol Catal 1996, V111, P267 HCAPLUS  
 (33) Vogt, D; Applied Homogenous Catalysis with Organometallic Compounds 1996, V1, P245 HCAPLUS  
 (34) Wang, L; J Am Chem Soc 1992, V114, P3169 HCAPLUS  
 (35) Wilke, G; Angew Chem Int Ed Engl 1988, V27, P185  
 (36) Ziegler, K; Angew Chem 1955, V67, P541 HCAPLUS  
 IT 204203-10-1, 2,6-Bis[1-(2,6-diisopropylphenylimino)ethyl]pyridinei  
 ron(II) chloride 207129-97-3

RL: CAT (Catalyst use); USES (Uses)

--(highly active iron and cobalt catalysts with bulky pyridinebisimine  
 ligands for olefin polymn.)


RN 204203-10-1 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliidyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)



RN 207129-97-3 HCAPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyliidyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]]-, (SP-5-13)- (9CI) (CA INDEX NAME)



DN 131:214605  
 TI Oligomerization of Olefins Catalyzed by New Cationic Palladium(II)  
 Complexes Containing an Unsymmetrical  $\alpha$ -Diimine Ligand  
 AU Meneghetti, Simoni Plentz; Lutz, Pierre J.; Kress, Jacky  
 CS Institut Charles Sadron Laboratoire de Chimie des Metaux de Transition et  
 de Catalyse Institut Le Bel, UPR 22 du CNRS UMR 7513 du CNRS Universite  
 Louis Pasteur, Strasbourg, 67000, Fr.  
 SO Organometallics (1999), 18(15), 2734-2737  
 CODEN: ORGND7; ISSN: 0276-7333  
 PB American Chemical Society  
 DT Journal  
 LA English  
 CC 35-3 (Chemistry of Synthetic High Polymers)  
 Section cross-reference(s): 29, 75  
 AB The palladium complexes  $[(Py-2-CMe:NAr)PdMe(MeCN)] + BAr'4-$  ( $2+BAr'4-$ ) and  
 $[Pd(Py-2-CMe:NAr)2]2+(BAr'4-)2$  [ $Ar = 2,6-(i-Pr)2-C6H3$ ,  $Ar' =$   
 $3,5-(CF3)2-C6H3$ )] have been synthesized from  $(Py-2-CMe:NAr)PdMeCl$ , and  
 their crystal and mol. structure has been detd. by X-ray anal.  $2+BAr'4-$   
 is an efficient catalyst for the oligomerization of ethylene, propylene,  
 and 1-hexene, as well as for the co-oligomerization of ethylene with alkyl  
 acrylates. P P P P P P 175273-54-8P RL: SPN (Synthetic prepn.); PREP  
 (Prepn.) (prepn. of) IT 53346-06-8P 175083-61-1P 175083-62-2P 175273-51-5.  
 ST oligomerization alkene catalyst palladium complex diimine; ethylene  
 oligomerization catalyst palladium complex diimine; propylene  
 oligomerization catalyst palladium complex diimine; hexene oligomerization  
 catalyst palladium complex diimine; crystal structure palladium complex  
 acetylidiisopropylphenylimine catalyst  
 IT Bond angle  
 Bond length  
 Crystal structure  
 (of catalysts; oligomerization of olefins catalyzed by cationic  
 palladium(II) complexes contg. unsym. (acetylidiisopropylphenylimine)pyr  
 idine ligand)  
 IT Molecular structure  
 (of catalysts; oligomerization of olefins catalyzed by cationic  
 palladium(II) complexes contg. unsym. (acetylidiisopropylphenylimine)pyr  
 idine ligand)  
 IT Polymerization catalysts  
 (oligomerization; oligomerization of olefins catalyzed by  
 cationic palladium(II) complexes contg. unsym.  
 (acetylidiisopropylphenylimine)pyridine ligand)  
 IT 63936-85-6 79060-88-1 219325-26-5  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (in catalyst prepn.; oligomerization of olefins catalyzed by cationic  
 palladium(II) complexes contg. unsym. (acetylidiisopropylphenylimine)pyr  
 idine ligand)  
 IT 242489-75-4P 242489-77-6P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (in catalyst prepn.; oligomerization of olefins catalyzed by cationic  
 palladium(II) complexes contg. unsym. (acetylidiisopropylphenylimine)pyr  
 idine ligand)  
 IT 9002-88-4P, Ethylene homopolymer 9003-07-0P 25067-06-5P, 1-Hexene  
 homopolymer 25101-13-7P, Ethylene-methyl methacrylate copolymer  
 104468-97-5P, Benzyl methacrylate-ethylene copolymer  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (oligomeric; oligomerization of olefins catalyzed by cationic  
 palladium(II) complexes contg. unsym. (acetylidiisopropylphenylimine)pyr  
 idine ligand)  
 IT 242791-16-8

RL: CAT (Catalyst use); USES (Uses)  
 (oligomerization of olefins catalyzed by cationic palladium(II)  
 complexes contg. unsym. (acetylidiisopropylphenylimine)pyridine  
 ligand)

IT 242489-79-8P

RL: SPN (Synthetic preparation); PREP (Preparation)  
 (oligomerization of olefins catalyzed by cationic palladium(II)  
 complexes contg. unsym. (acetylidiisopropylphenylimine)pyridine ligand)

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE


- (1) Bahr, S; J Org Chem 1992, V57, P5545 HCPLUS
- (2) Brookhart, M; J Am Chem Soc 1992, V114, P5894 HCPLUS
- (3) Chatt, L; J Chem Soc 1957, P34
- (4) Desjardin, S; Organometallics 1998, V17, P2382
- (5) Drent, E; Pure Appl Chem 1990, V62, P661 HCPLUS
- (6) Feldman, J; Organometallics 1997, V16, P1514 HCPLUS
- (7) Johnson, L; J Am Chem Soc 1995, V117, P6414 HCPLUS
- (8) Mecking, S; J Am Chem Soc 1998, V120, P888 HCPLUS
- (9) Rix, F; J Am Chem Soc 1995, V117, P1137 HCPLUS
- (10) Rulke, R; Inorg Chem 1993, V32, P5769 HCPLUS
- (11) Rulke, R; J Organomet Chem 1996, V508, P109

IT 242489-75-4P 242489-77-6P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (in catalyst prepn.; oligomerization of olefins catalyzed by cationic palladium(II) complexes contg. unsym. (acetylidiisopropylphenylimine)pyridine ligand)

RN 242489-75-4 HCPLUS

CN Palladium, [2,6-bis(1-methylethyl)-N-[1-(2-pyridinyl-.kappa.N)ethylidene]benzenamine-.kappa.N]chloromethyl-, (SP-4-3)- (9CI)  
 (CA INDEX NAME)




RN 242489-77-6 HCPLUS

CN Palladium(1+), (acetonitrile)[2,6-bis(1-methylethyl)-N-[1-(2-pyridinyl-.kappa.N)ethylidene]benzenamine-.kappa.N]methyl-, (SP-4-3)-, tetrakis[3,5-bis(trifluoromethyl)phenyl]borate(1-) (9CI) (CA INDEX NAME)


CM 1

CRN 242489-76-5  
 CMF C22 H30 N3 Pd  
 CCI CCS  
 CDES 7:SP-4-3



CM 2

CRN 79230-20-9  
 CMF C32 H12 B F24  
 CCI CCS



IT 242791-16-8


RL: CAT (Catalyst use); USES (Uses)  
 (oligomerization of olefins catalyzed by cationic palladium(II)  
 complexes contg. unsym. (acetylidiisopropylphenylimine)pyridine  
 ligand)

RN 242791-16-8 HCPLUS

CN Palladium(1+), (acetonitrile)[2,6-bis(1-methylethyl)-N-[1-(2-pyridinyl-.kappa.N)ethylidene]benzenamine-.kappa.N]methyl-, (SP-4-2)-,  
 tetrakis[3,5-bis(trifluoromethyl)phenyl]borate(1-) (9CI) (CA INDEX NAME)


CM 1

CRN 242791-15-7  
 CMF C22 H30 N3 Pd  
 CCI CCS  
 CDES 7:SP-4-2



CM 2

CRN 79230-20-9  
 CMF C32 H12 B F24  
 CCI CCS



L31 ANSWER 16 OF 17 HCPLUS COPYRIGHT 2002 ACS  
 AN 1998:723809 HCPLUS  
 DN 130:4185  
 TI Metal complexes having nitrogen-containing ligands as **olefin polymerization catalysts**  
 IN Bres, Philippe-Luc; Gibson, Vernon Charles; Mabille, Christine Daniele Florence; Reed, Warren; Wass, Duncan; Weatherhead, Richard Henry  
 PA BP Chemicals Ltd., UK  
 SO PCT Int. Appl., 25 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C08F010-00  
 ICS C08F004-70; C08F004-80; C07F015-00; C07F015-04  
 CC 35-3 (Chemistry of Synthetic High Polymers)  
 Section cross-reference(s): 29

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9849208	A1	19981105	WO 1998-GB1205	19980424
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG

AU 9870693 A1 19981124 AU 1998-70693 19980424

PRAI GB 1997-8487 19970425  
WO 1998-GB1205 19980424

OS MARPAT 130:4185  
GI



AB A nitrogen-contg. metal complex compd. I wherein the ring formed by Q, Cl<sub>1</sub>, and N<sub>1</sub> is aromatically unsatd., the divalent group Q comprises a chain of 3-4 atoms having the formula (CR)<sub>n</sub>(Z)<sub>r</sub> [n = 2, 3, 4; r = 0, 1; (n + r) = 3, 4; Z = N, O, S; R = H, halogen, C1-30 hydrocarbyl, NR42, OR5, NO2; R<sub>4</sub>, R<sub>5</sub> = C1-30 hydrocarbyl], R<sub>1</sub> is H, halogen, or C1-30 hydrocarbyl or halohydrocarbyl, R<sub>2</sub> is H, halogen, or C1-30 hydrocarbyl, M is nickel or palladium, X is a univalent radical, for example, halide, hydride, hydrocarbyl oxide.. Q is preferably a pyridine ring. Also described is a catalyst for the **polymn.** of 1-**olefins** comprising the defined metal complex and an activating quantity of a compd. selected from organoaluminium compds., aluminoxanes and fluorohydrocarbylboron compds.

ST **polymn olefin** catalyst metal complex; palladium complex nitrogen ligand polymn catalyst; nickel complex nitrogen ligand polymn catalyst

IT Aluminoxanes  
RL: CAT (Catalyst use); USES (Uses)  
(Me; metal complexes having nitrogen-contg. ligands as **olefin polymn. catalysts**)

IT Polymerization  
(gas-phase; metal complexes having nitrogen-contg. ligands as **olefin polymn. catalysts**)

IT Polymerization catalysts  
(metal complexes having nitrogen-contg. ligands as **olefin polymn. catalysts**)

IT Polymerization  
(slurry; metal complexes having nitrogen-contg. ligands as **olefin polymn. catalysts**)

IT Polymerization  
(soln.; metal complexes having nitrogen-contg. ligands as **olefin polymn. catalysts**)

IT 1314-23-4, Zirconia, uses 1344-28-1, Alumina, uses 7631-86-9, Silica, uses 79060-88-1 139362-04-2  
RL: CAT (Catalyst use); USES (Uses)  
(metal complexes having nitrogen-contg. ligands as **olefin polymn. catalysts**)

IT 215594-94-8P 215594-96-0P 215594-97-1P  
215594-99-3P 215595-01-0P 215595-03-2P

215595-05-4P

RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);  
 USES. (Uses)

(metal complexes having nitrogen-contg. ligands as  
 olefin polymn. catalysts)

IT 9002-88-4P

RL: IMF (Industrial manufacture); PREP (Preparation)  
 (metal complexes having nitrogen-contg. ligands as olefin  
 polymn. catalysts)

IT 5780-66-5, Pyrazinecarboxaldehyde 24544-04-5, 2,6-Diisopropylaniline  
 28923-39-9 63936-85-6 149810-35-5 215459-01-1

RL: RCT (Reactant); RACT (Reactant or reagent)

(prepn. of metal complexes having nitrogen-contg. ligands as  
 olefin polymn. catalysts)

IT 215459-03-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (prepn. of metal complexes having nitrogen-contg. ligands as  
 olefin polymn. catalysts)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Canovese, L; JOURNAL OF ORGANOMETALLIC CHEMISTRY 1996, V508(1), P101
- (2) de Felice, V; JOURNAL OF ORGANOMETALLIC CHEMISTRY 1995, V493(1), P1
- (3) Du Pont; WO 9702298 A 1997 HCAPLUS
- (4) Orrell, K; JOURNAL OF ORGANOMETALLIC CHEMISTRY 1997, V530(1-2), P235  
 HCAPLUS
- (5) Rulke, R; JOURNAL OF ORGANOMETALLIC CHEMISTRY 1996, V508(1), P109


IT 215594-94-8P 215594-96-0P 215594-97-1P  
 215594-99-3P 215595-01-0P 215595-03-2P

215595-05-4P

RL: CAT (Catalyst use); SPN (Synthetic preparation); PREP (Preparation);  
 USES (Uses)


(metal complexes having nitrogen-contg. ligands as  
 olefin polymn. catalysts)

RN 215594-94-8 HCAPLUS

CN Nickel, [2,6-bis(1-methylethyl)-N-[(2-pyridinyl-  
 .kappa.N)methylene]benzenamine-.kappa.N]dibromo- (9CI) (CA INDEX NAME)


RN 215594-96-0 HCAPLUS

CN Nickel, [2,6-bis(1-methylethyl)-N-[(6-methyl-2-pyridinyl-  
 .kappa.N)methylene]benzenamine-.kappa.N]dibromo- (9CI) (CA INDEX NAME)




RN 215594-97-1 HCAPLUS

CN Nickel, [2,6-bis(1-methylethyl)-N-[(2-pyridinyl-.kappa.N)methylene]benzenamine-.kappa.N]dimethyl- (9CI) (CA INDEX NAME)




RN 215594-99-3 HCAPLUS

CN Palladium, [2,6-bis(1-methylethyl)-N-[(2-pyridinyl-.kappa.N)methylene]benzenamine-.kappa.N]chloromethyl- (9CI) (CA INDEX NAME)




RN 215595-01-0 HCAPLUS

CN Palladium, [2,6-bis(1-methylethyl)-N-[(2-pyridinyl-.kappa.N)methylene]benzenamine-.kappa.N]dimethyl- (9CI) (CA INDEX NAME)




RN 215595-03-2 HCAPLUS

CN Nickel, [2,6-bis(1-methylethyl)-N-[(pyrazinyl-.kappa.N1)methylene]benzenamine-.kappa.N]dibromo- (9CI) (CA INDEX NAME)



RN 215595-05-4 HCAPLUS

CN Nickel, [2,6-bis(1-methylethyl)-N-[(pyrazinyl-.kappa.N1)methylene]benzenamine-.kappa.N]dimethyl- (9CI) (CA INDEX NAME)



L31 ANSWER 17 OF 17 HCAPLUS COPYRIGHT 2002 ACS

AN 1998:270894 HCAPLUS

DN 128:321993

TI Novel olefin polymerization catalysts based on iron and cobalt

AU Birtovsek, George J. P.; Gibson, Vernon; Kimberley, Brian S.; Maddox, Peter J.; McTavish, Stuart J.; Solan, Gregory A.; White, Andrew J. P.; Williams, David J.

CS Dep. Chem., Imperial Coll., Kensington, London, SW7 2AY, UK

SO Chemical Communications (Cambridge) (1998), (7), 849-850

CODEN: CHCOFS; ISSN: 1359-7345

PB Royal Society of Chemistry

DT Journal

LA English

CC 35-3 (Chemistry of Synthetic High Polymers)

AB Catalysts derived from iron and cobalt complexes bearing 2,6-bis(imino)pyridyl ligands for polymn. of ethylene were studied. All of the catalysts produced essentially linear polyethylene with mol. wts. that were dependent upon the aryl substitution pattern. There was a marked dependence of the polymer mol. wt. on ethylene pressure for the iron catalyst system. The cobalt catalyst displayed a considerably lower activity than its iron analog.

ST iron iminopyridyl polymn catalyst ethylene; cobalt iminopyridyl polymn catalyst ethylene; polyethylene prep catalyst iron cobalt

IT Polymerization catalysts

(iron and cobalt complexes contg. bis(imino)pyridyl ligands as catalysts for polymn. of ethylene)

IT 204203-10-1 207129-93-9 207129-94-0

207129-95-1 207129-96-2 207129-97-3

RL: CAT (Catalyst use); USES (Uses)

(iron and cobalt complexes contg. bis(imino)pyridyl ligands as catalysts for polymn. of ethylene)

IT 9002-88-4P, Polyethylene

RL: SPN (Synthetic preparation); PREP (Preparation)

(iron and cobalt complexes contg. bis(imino)pyridyl ligands as catalysts for polymn. of ethylene)

IT 204203-10-1 207129-93-9 207129-94-0


207129-95-1 207129-96-2 207129-97-3

RL: CAT (Catalyst use); USES (Uses)

(iron and cobalt complexes contg. bis(imino)pyridyl ligands as catalysts for polymn. of ethylene)


RN 204203-10-1 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyli dyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]-, (SP-5-13)- (9CI) (CA INDEX NAME)




RN 207129-93-9 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyli dyne]bis[2,6-dimethylbenzenamine-.kappa.N]-, (SP-5-13)- (9CI) (CA INDEX NAME)




RN 207129-94-0 HCPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyli dyne]bis[2,4,6-trimethylbenzenamine-.kappa.N]-, (TB-5-22)- (9CI) (CA INDEX NAME)




RN 207129-95-1 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethylidyne]bis[2,4-dimethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



RN 207129-96-2 HCAPLUS

CN Iron, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)dimethylidyne]bis[2,6-dimethylbenzenamine-.kappa.N]- (9CI) (CA INDEX NAME)



RN 207129-97-3 HCAPLUS

CN Cobalt, dichloro[N,N'-(2,6-pyridinediyl-.kappa.N)diethyldyne]bis[2,6-bis(1-methylethyl)benzenamine-.kappa.N]-, (SP-5-13)- (9CI) (CA INDEX NAME)

